


# Love takes up where pain leaves off, brain study shows

October 13 2010

---



Love-induced pain relief was associated with the activation of primitive brain structures that control rewarding experiences, such as the nucleus accumbens - shown here in color. Courtesy of Sean Mackey and Jarred Younger

(PhysOrg.com) -- Intense, passionate feelings of love can provide amazingly effective pain relief, similar to painkillers or such illicit drugs as cocaine, according to a new Stanford University School of Medicine study.

"When people are in this passionate, all-consuming phase of [love](#), there are significant alterations in their mood that are impacting their experience of [pain](#)," said Sean Mackey, MD, PhD, chief of the Division of Pain Management, associate professor of anesthesia and senior author

of the study, which will be published online Oct. 13 in [PLoS ONE](#).

"We're beginning to tease apart some of these reward systems in the [brain](#) and how they influence pain. These are very deep, old systems in our brain that involve dopamine - a primary neurotransmitter that influences mood, reward and motivation."

Scientists aren't quite yet ready to tell patients with chronic pain to throw out the painkillers and replace them with a passionate love affair; rather, the hope is that a better understanding of these neural-rewards pathways that get triggered by love could lead to new methods for producing pain relief.

"It turns out that the areas of the brain activated by intense love are the same areas that drugs use to reduce pain," said Arthur Aron, PhD, a professor of psychology at State University of New York at Stony Brook and one of the study's authors. Aron has been studying love for 30 years. "When thinking about your beloved, there is intense activation in the reward area of the brain - the same area that lights up when you take cocaine, the same area that lights up when you win a lot of money."

The concept for the study was sparked several years ago at a neuroscience conference when Aron, an expert in the study of love, met up with Mackey, an expert in the research of pain, and they began talking.

"Art was talking about love," Mackey said. "I was talking about pain. He was talking about the brain systems involved with love. I was talking about the brain systems involved with pain. We realized there was this tremendous overlapping system. We started wondering, 'Is it possible that the two modulate each other?'"

After the conference, Mackey returned to Stanford and collaborated with postdoctoral scholar Jarred Younger, PhD, now an assistant

professor of [anesthesia](#), who was also intrigued with the idea. Together the three set up a study that would entail examining the brain images of undergraduates who claimed to be "in that first phase of intense love."

"We posted fliers around Stanford University and within hours we had undergrads banging on our door," Mackey said. The fliers asked for couples who were in the first nine months of a romantic relationship.

"It was clearly the easiest study the pain center at Stanford has ever recruited for," Mackey said. "When you're in love you want to tell everybody about it.

"We intentionally focused on this early phase of passionate love," he added. "We specifically were not looking for longer-lasting, more mature phases of the relationship. We wanted subjects who were feeling euphoric, energetic, obsessively thinking about their beloved, craving their presence.

"When passionate love is described like this, it in some ways sounds like an addiction. We thought, 'Maybe this does involve similar brain systems as those involved in addictions which are heavily dopamine-related.' Dopamine is the neurotransmitter in our brain that is intimately involved with feeling good."

Researchers recruited 15 undergraduates (eight women and seven men) for the study. Each was asked to bring in photos of their beloved and photos of an equally attractive acquaintance. The researchers then successively flashed the pictures before the subjects, while heating up a computer-controlled thermal stimulator placed in the palm of their hand to cause mild pain. At the same time, their brains were scanned in a functional magnetic resonance imaging machine.

The undergraduates were also tested for levels of pain relief while being

distracted with word-association tasks such as: "Think of sports that don't involve balls." Scientific evidence has shown in the past that distraction causes pain relief, and researchers wanted to make sure that love was not just working as a distraction from pain.

Results showed that both love and distraction did equally reduce pain, and at much higher levels than by concentrating on the photo of the attractive acquaintance, but interestingly the two methods of pain reduction used very different brain pathways.

"With the distraction test, the brain pathways leading to pain relief were mostly cognitive," Younger said. "The reduction of pain was associated with higher, cortical parts of the brain. Love-induced analgesia is much more associated with the reward centers. It appears to involve more primitive aspects of the brain, activating deep structures that may block pain at a spinal level - similar to how opioid analgesics work.

"One of the key sites for love-induced analgesia is the nucleus accumbens, a key reward addiction center for opioids, cocaine and other drugs of abuse. The region tells the brain that you really need to keep doing this," Younger said.

"This tells us that you don't have to just rely on drugs for [pain relief](#)," Aron said. "People are feeling intense rewards without the side effects of drugs."

Provided by Stanford University Medical Center

Citation: Love takes up where pain leaves off, brain study shows (2010, October 13) retrieved 24 November 2023 from <https://medicalxpress.com/news/2010-10-pain-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.