

Liver, belly fat may identify high risks of heart disease in obese people

July 21 2011

Obese people with high levels of abdominal fat and liver fat may face increased risks for heart disease and other serious health problems, according to research published in *Arteriosclerosis, Thrombosis and Vascular Biology: Journal of the American Heart Association*.

Obesity is commonly associated with heart disease risk and problems called cardiometabolic abnormalities, including [insulin resistance](#), [type 2 diabetes](#), cholesterol disorders, hypertension and gout.

Researchers in Sweden and Finland found that obese people at the highest risk have increased secretion of liver lipids, more abdominal fat and impaired removal of triglycerides from the blood stream. As such, doctors should routinely check [obese patients](#) for intra-abdominal obesity and indications of liver fat, researchers said.

"It is important to recognize that measuring abdominal fat and liver fat can identify the patients at high risk for [metabolic abnormalities](#) and heart disease," said Jan Boren, M.D., Ph.D., study senior author and professor of molecular and clinical medicine at the University of Gothenburg in Sweden. "Such exams are important because up to 20 percent of the obese appear to be "metabolically normal."

The researchers investigated what makes some obese people develop lipid disorders. They found that liver fat is strongly associated with increased secretion of very-low-density lipoproteins (VLDL), which contain the highest amount of triglycerides. High levels of triglycerides

carry an increased risk of metabolic abnormalities and increased risk of heart disease and premature death.

"Increased liver fat is dangerous, as it is linked with many known [heart disease risk](#) factors," said Marja-Riitta Taskinen, M.D., Ph.D. study co-author and professor of medicine at the University of Helsinki in Finland. "[Lifestyle modifications](#) such as exercise and weight loss can reduce liver fat and the secretion of lipoproteins."

In the study, middle-aged Caucasian Finnish men were divided into three groups: 14 obese men had high triglycerides; 14 obese men had normal triglycerides and 10 men who were not obese and had normal triglycerides served as controls. The obese men's average age was 52 to 55 years. The non-obese men's average age was 48 years.

Researchers used proton magnetic resonance spectroscopy to measure liver fat, while subcutaneous abdominal and visceral fat were measured by magnetic resonance imaging.

The results showed that triglyceride levels in the obese men were higher as a result of the increased secretion coupled with severely impaired clearance of triglyceride-rich VLDL particles. Researchers also found:

- The obese men with elevated triglycerides had a liver fat content of 13 percent.
- The obese men with normal triglycerides had a 6.9 percent liver fat content.
- The non-obese men had a 2.9 percent liver fat content.

Researchers plan to conduct a similar study in women.

Provided by American Heart Association

Citation: Liver, belly fat may identify high risks of heart disease in obese people (2011, July 21)
retrieved 20 November 2023 from

<https://medicalxpress.com/news/2011-07-liver-belly-fat-high-heart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.