

Incontinence 20 years after child birth three times more common after vaginal delivery

March 26 2012

Women are nearly three times more likely to experience urinary incontinence for more than 10 years following a vaginal delivery rather than a caesarean section, finds new research at the Sahlgrenska Academy at the University of Gothenburg, Sweden.

Urinary incontinence (UI) is a common condition affecting [adult women](#) of all ages and can have a [negative influence](#) on quality of life.

This new study from the University of Gothenburg in Sweden looked at the prevalence and risk factors for UI 20 years after vaginal delivery (VD) or [caesarean section](#) (CS). The study included women who had only one child and assessed their prevalence of UI for less than five years, between 5-10 years and for more than 10 years.

The SWEPOP (Swedish pregnancy, obesity and pelvic floor) study was conducted in 2008 and data were obtained from the Medical Birth Register (MBR) for deliveries between 1985 and 1988. A questionnaire was sent to women and 6, 148 completed it answering questions on height, weight, urinary or anal incontinence, genital prolapse, menstrual status, hysterectomy, the menopause and [hormone treatment](#).

Overall, the prevalence of UI was considerably higher after a vaginal delivery (40.3%) compared to women who delivered by caesarean section (28.8%).

The study also found that the prevalence of UI for more than 10 years

almost tripled after VD (10.1%) compared to women who had a CS (3.9%).

In addition, the paper looks at the impact of BMI on UI. The risk increase of UI in obese women more than doubled in comparison to women with a normal BMI after VD and more than tripled after CS.

Maria Gyhagen, Department of [Obstetrics and Gynaecology](#), Sahlgrenska Academy, University of Gothenburg, Sweden and co-author of the paper said: “In conclusion, the risk of developing [urinary incontinence](#) was higher 20 years after a vaginal delivery compared to a caesarean section. “There are many factors affecting urinary incontinence but obesity and ageing as well as obstetric trauma during childbirth are known to be three of the most important risk factors.”

BJOG Deputy Editor-in-Chief, Pierre Martin-Hirsch, added: “Urinary incontinence affects many women and can have a big impact on day to day life. “However, women need to look at all the information when deciding on mode of delivery as despite [vaginal delivery](#) and BMI being linked to urinary incontinence, caesarean section involves its own risks.”

Provided by University of Gothenburg

Citation: Incontinence 20 years after child birth three times more common after vaginal delivery (2012, March 26) retrieved 2 February 2024 from <https://medicalxpress.com/news/2012-03-incontinence-years-child-birth-common.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--