

## Study examines use of waist measures among overweight and obese adolescents

April 2 2012

---

Waist measures (waist circumference, waist to height ratio) in conjunction with body mass index appear to be associated with lipid and blood pressure assessments among overweight and obese adolescents, according to a report published Online First by *Archives of Pediatrics & Adolescent Medicine*.

Body mass index (BMI) is currently the most commonly used measure of obesity in adolescents, according to background in the study. BMI cannot differentiate between fat and fat-free [body mass](#) however, so the authors included additional obesity measures including [waist circumference](#) (WC) and waist to height ratio (WHtR; WC divided by height) in relation to BMI to examine the role of waist measures in the [lipid](#) and [blood pressure](#) assessment of adolescents classified by BMI.

Michael Khoury, M.D., of the University of Toronto, and the Hospital for Sick Children, Ontario, Canada, and colleagues, analyzed data collected on a population of 14- and 15-year-old students in the Niagara Region, Ontario. Data were collected during the students' mandatory physical education class and a total of 4,104 students participated, with 3,248 students submitting full data for inclusion in the analysis.

Overall, the authors found that the association between blood pressure, lipid profile, and measures of obesity (BMI alone, BMI/WC percentile and BMI/WHtR) were statistically significant, but the associations had limited strength and were similar using each measure of obesity. After adjusting for the child's age and sex, among overweight and obese

adolescents, increasing WHtR categories corresponded with worsening lipid profile, with the greatest associations occurring in obese adolescents.

Obese adolescents also had increased odds of a higher degree of hypertension, and obese adolescents with increased WHtR had further increased odds of hypertension compared to those adolescents with more normal waist measures.

"In conclusion, waist measures serve to further specify lipid and blood pressure assessments in overweight and obese adolescents, with the greatest associations noted for obese adolescents," the authors write.

"Waist measures appear to be important discriminating measurements when assessing lipid and blood pressure measurements in adolescents with high BMI and should be included when screening for cardiometabolic risk in overweight and obese [adolescents](#)."

**More information:** *Arch Pediatr Adolesc Med*. Published online April 2, 2012. [doi:10.1001/archpediatrics.2012.126](https://doi.org/10.1001/archpediatrics.2012.126)

Provided by JAMA and Archives Journals

Citation: Study examines use of waist measures among overweight and obese adolescents (2012, April 2) retrieved 20 November 2023 from <https://medicalxpress.com/news/2012-04-waist-overweight-obese-adolescents.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--