

Inducing labor can reduce birth complications without increasing Cesareans

May 10 2012

Induction of labour beyond 37 weeks of pregnancy can reduce perinatal mortality (death before, during or shortly after birth) without increasing caesarean section rates, finds a study published on *BMJ* today.

However, induction is associated with a greater risk of admission to a special care baby unit.

Induction of labour is frequently performed for pregnancies of more than 41 weeks gestation to reduce perinatal mortality. Induction at term (beyond 37 weeks) can also reduce complications, particularly when a mother has existing health problems like [high blood pressure](#).

Yet there has been no large study examining the risks and benefits of induction at term on [newborn deaths](#).

This information is needed to help guide decisions about [pregnancy](#) management. So a team of researchers in Scotland compared rates of perinatal mortality and maternal complications after elective induction (induction of labour with no recognised medical indication) and expectant management (continuation of pregnancy to either spontaneous labour or induction or [caesarean section](#) at a later date).

Using Scottish birth and death records, they analysed data for over 1.2 million women with single pregnancies who gave birth after 37 weeks gestation between 1981 and 2007. Outcomes were adjusted for factors such as age at delivery, whether the mother had given birth before, [birth](#)

[weight](#) and deprivation.

At each gestation between 37 and 41 weeks, elective induction of labour was associated with lower [death rates](#) compared with expectant management, without increasing the need for a caesarean section. For example, at 40 weeks gestation, deaths occurred in 37 out of 44,764 (0.08%) in the induction group compared with 627 out of 350,643 (0.18%) in the expectant management group.

However, elective induction of labour was associated with increased rates of admission to a special care baby unit (8%) compared with expectant management (7.3%).

This means that for every 1,040 women having elective induction of labour at 40 weeks, one newborn death may be prevented, but this would result in seven more admissions to a special care baby unit, explain the authors.

Although there's a possibility that some other unmeasured (confounding) factor may explain these results, the authors conclude that elective induction of labour at term "has the potential to reduce [perinatal mortality](#) in developed countries without increasing the risk of operative delivery."

Provided by British Medical Journal

Citation: Inducing labor can reduce birth complications without increasing Cesareans (2012, May 10) retrieved 19 November 2023 from <https://medicalxpress.com/news/2012-05-labor-birth-complications-cesareans.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.