

Filaggrin mutations up risk of irritant contact dermatitis

October 14 2012



Image courtesy of Blausen Medical

Both atopic dermatitis and loss-of-function mutations in the filaggrin gene are independently associated with an increased risk of developing chronic irritant contact dermatitis, with people having both mutations at about a five-fold higher risk, according to research published online Oct. 5 in the *British Journal of Dermatology*.

(HealthDay)—Both atopic dermatitis (AD) and loss-of-function mutations in the filaggrin gene (*FLG*) are independently associated with an increased risk of developing chronic irritant contact dermatitis (ICD), with people having both mutations at about a five-fold higher risk, according to research published online Oct. 5 in the *British Journal of Dermatology*.

Noting that loss-of-function mutations in *FLG* increase the risk for AD, Maaïke J. Visser, M.D., of the Coronel Institute for Occupational Health in Amsterdam, and colleagues conducted a study involving 634 subjects with chronic ICD and 393 controls to investigate the relative

contribution and interaction of *FLG* mutations and AD in ICD.

The researchers found that 15.9 percent of ICD patients and 8.3 percent of control patients had an *FLG* mutation, with a crude odds ratio of 2.09 for the combined carrier allele. After correcting for AD, the adjusted odds ratio for *FLG* mutations was 1.62, and individuals with AD had an odds ratio of 2.89 for developing ICD. Concomitant presence of both AD and *FLG* mutations resulted in a 4.7-fold increased risk of ICD.

"In summary, our results indicate that both *FLG* loss-of-function mutations and AD significantly increase the risk for ICD, with respective odds ratios of 1.61 and 2.89," the authors write. "Individuals with both *FLG* [mutations](#) and AD have an approximately four- to five-fold increased risk to develop ICD."

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2012 [HealthDay](#). All rights reserved.

Citation: Filaggrin mutations up risk of irritant contact dermatitis (2012, October 14) retrieved 22 November 2023 from <https://medicalxpress.com/news/2012-10-filaggrin-mutations-contact-dermatitis.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--