

Pulmonary hypertension combination therapy may lead to greater disease burden

October 22 2012

Patients with pulmonary arterial hypertension (PAH) receiving combination therapy with intravenous (IV) PGI₂ may suffer from greater disease burden compared with those receiving monotherapy or combination therapy, excluding IV PGI₂.

Researchers from Massachusetts General Hospital and Novartis Pharma AG in Switzerland performed statistical tests on 446 patients with PAH, of which (a) 337 were receiving monotherapy; (b) 88 were receiving non-IV PGI₂ combination therapy; and (c) 21 were receiving IV PGI₂ [combination therapy](#).

Results showed that pulmonary vascular resistance, mean number of PAH-related hospitalizations, and dyspnea scores were highest in those treated with IV PGI₂.

This study was presented during CHEST 2012, the annual meeting of the [American College of Chest Physicians](#), held October 20 – 25, in Atlanta, Georgia.

Provided by American College of Chest Physicians

Citation: Pulmonary hypertension combination therapy may lead to greater disease burden (2012, October 22) retrieved 4 February 2024 from <https://medicalxpress.com/news/2012-10-pulmonary-hypertension-combination-therapy-greater.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.