


HIV-HCV coinfection speeds HCV-related liver fibrosis

February 27 2013



Individuals who are coinfectd with HIV and hepatitis C virus exhibit liver fibrosis similar to that of individuals without HIV who are nearly 10 years older, according to research published online Feb. 26 in the *Annals of Internal Medicine*.

(HealthDay)—Individuals who are coinfectd with HIV and hepatitis C virus (HCV) exhibit liver fibrosis similar to that of individuals without HIV who are nearly 10 years older, according to research published online Feb. 26 in the *Annals of Internal Medicine*.

Gregory D. Kirk, M.D., M.P.H., Ph.D., of Johns Hopkins Bloomberg School of Public Health in Baltimore, and colleagues conducted a cohort study involving 1,176 current and former injection drug users with antibodies to HCV (median age, 49 years) to examine whether individuals with concurrent HIV infection develop HCV-related liver disease at younger ages than those without HIV infection.

Overall, 34 percent of participants were coinfecting with HIV and HCV. Compared with patients who only had HCV, the researchers found that those who were coinfecting had a significantly higher prevalence of clinically significant fibrosis without cirrhosis and of cirrhosis.

Independent correlates of [liver fibrosis](#) included increasing age and HIV infection, as well as daily alcohol use, chronic [hepatitis B virus](#) infection, [body mass index](#) greater than 25 kg/m², and elevated plasma HCV RNA levels. After accounting for these factors, HIV-infected patients exhibited liver fibrosis equal to that of people without HIV who were an average of 9.2 years older.

"In summary, persons who are coinfecting with HIV and HCV have liver fibrosis stages similar to persons infected with HCV alone who are nearly one decade older," the authors write. "Additional work is needed to understand the biological basis for this observation and investigate whether effective treatment of HIV and HCV infections reduce future liver fibrosis progression."

One of the authors disclosed financial ties to the pharmaceutical industry. Support of reagents for HCV RNA testing was provided by Abbott Molecular.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

[Health News](#) Copyright © 2013 [HealthDay](#). All rights reserved.

Citation: HIV-HCV coinfection speeds HCV-related liver fibrosis (2013, February 27) retrieved 20 November 2023 from <https://medicalxpress.com/news/2013-02-hiv-hcv-coinfection-hcv-related-liver-fibrosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.