

Stillbirth rates have increased significantly, although spontaneous stillbirth rates have not

April 8 2013

The rate of stillbirths in British Columbia, Canada, increased by 31% over a decade, although the rate of spontaneous stillbirths did not increase, according to a study published in *CMAJ (Canadian Medical Association Journal)*.

Stillbirth rates, which had been declining for decades, have increased or plateaued in several [industrialized countries](#). For example, rates in Australia have increased from 7.0 per 1000 total births in 2000 to 7.8 per 1000 in 2009 and in Canada, rates have increased from 6.0 per 1000 total births in 2000 to 7.1 per 1000 in 2009. The United States, United Kingdom and New Zealand have also seen rates increase or plateau.

"The prevention of stillbirth through early delivery of the compromised fetus is considered the cornerstone of modern obstetrics," writes Dr. K.S. Joseph, Department of [Obstetrics and Gynaecology](#) and School of Population and Public Health, University of British Columbia, with coauthors. "The recent trends in stillbirth rates are therefore disquieting given developments in fetal surveillance techniques, increased monitoring of high-risk pregnancies and simultaneous increases in obstetric intervention."

To understand the reasons for these recent increases, BC researchers looked at data on all stillbirths of 20 weeks' gestation or older in the province from 2000 to 2010. There were 461 083 live births and 3991 stillbirths in the 10-year period. The rate of stillbirths increased 31%, from 8.08 per 1000 total births in 2000 to 10.55 per 1000 in 2010.

Stillbirths among [fetuses](#) under 500 g [birth weight](#) increased significantly, from 2.36 per 1000 total births to 2.84 per 1000 but decreased from 2.73 to 1.95 per 1000 total births in the 1000-g or heavier category. Stillbirths before 22 weeks' gestation increased from 2.71 per 1000 total births to 3.71 per 1000. The study found that the increase in stillbirth rates was due to increases in [prenatal diagnosis](#) and [pregnancy termination](#) of fetuses with congenital anomalies or other reasons.

"Increases in pregnancy terminations were responsible for the increases observed in stillbirth rates from 2000 to 2010 and were associated with declines in the prevalence of congenital abnormalities among live-born infants," write the authors.

After excluding stillbirths due to pregnancy terminations, the researchers found a non-significant decline in the rate of spontaneous stillbirths.

They suggest that "advances in, and increased uptake of, prenatal diagnosis are having a population-level impact on the timing of death and the gestational age at delivery of fetuses with congenital anomalies."

On the other hand, the researchers identified archaic stillbirth registration rules that place an undue psychosocial burden on mothers and families. "Legal requirements for stillbirth registration add unnecessary stress to grieving parents and should be revamped," says Cheryl Davies, an author of the study and Vice-President, Ambulatory Programs at the Women's Hospital and Health Centre of British Columbia.

"Future research examining temporal trends and regional variations in [stillbirth](#) rates should focus separately on spontaneous stillbirths and those due to pregnancy termination," they conclude.

More information: www.cmaj.ca/lookup/doi/10.1503/cmaj.121372

Provided by Canadian Medical Association Journal

Citation: Stillbirth rates have increased significantly, although spontaneous stillbirth rates have not (2013, April 8) retrieved 18 May 2023 from

<https://medicalxpress.com/news/2013-04-stillbirth-significantly-spontaneous.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.