

# Delayed transfer to the ICU increases risk of death in hospital patients

May 21 2013

---

Delayed transfer to the intensive care unit (ICU) in hospitalized patients significantly increases the risk of dying in the hospital, according to a new study from researchers in Chicago.

"[Early intervention](#) improves outcomes for many of the conditions that are indications for inpatient transfer to the ICU. This suggests that delaying ICU transfer may increase the risk of death in these patients," said lead author Matthew Churpek, MD, MPH, of the University of Chicago Medical Center. "Using a vital sign-based early warning score, the [Cardiac Arrest](#) Risk Triage (CART) score, we examined outcomes after delayed transfer to the ICU and found that each one hour increase in transfer delay was associated with a 7% increase in the odds of dying in the ICU."

The results of the study will be presented at the ATS 2013 International Conference.

The study included 2,166 patients at an [academic hospital](#) who were transferred from medical-surgical wards to the ICU. Of these patients, 425 (20%) died during admission.

In a subset of 260 patients transferred to the ICU within 6 hours of a first critical CART score value, 71(27%) died during [hospital admission](#), compared with 65 of the 143 (45%) patients for whom ICU transfer was delayed for more than 6 hours(p

Citation: Delayed transfer to the ICU increases risk of death in hospital patients (2013, May 21)  
retrieved 20 November 2023 from

<https://medicalxpress.com/news/2013-05-icu-death-hospital-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.