

Understanding the molecular mechanisms underlying Alzheimer's disease

June 10 2013

The accumulation of amyloid- β ($A\beta$) in the brains of Alzheimer's disease (AD) patients is known to be associated with memory loss and neuronal degeneration, but the mechanism of $A\beta$ pathogenesis is not fully understood.

In this issue of the *Journal of Clinical Investigation*, researchers led by Yong-Keun Jung at Seoul National University demonstrate that $A\beta$ binds to a [cellular protein](#) known as FC γ RIIb.

Greater levels of FC γ RIIb were detected in the brains of AD patients. Binding of $A\beta$ to FC γ RIIb activated cell stress and death pathways. In a mouse model of AD, depletion of FC γ RIIb ameliorated memory impairment.

This study demonstrates that FC γ RIIb plays a critical role in AD pathogenesis.

More information: Fc γ RIIb mediates amyloid- β neurotoxicity and memory impairment in Alzheimer's disease, *J Clin Invest*.

[doi:10.1172/JCI66827](https://doi.org/10.1172/JCI66827)

Provided by Journal of Clinical Investigation

Citation: Understanding the molecular mechanisms underlying Alzheimer's disease (2013, June

10) retrieved 19 November 2023 from <https://medicalxpress.com/news/2013-06-molecular-mechanisms-underlying-alzheimer-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.