

## Marked variability seen in total knee arthroplasty indications

June 14 2013



There is considerable variability associated with total knee arthroplasty recommendations, according to a report published in the May issue of *Pain Medicine News*.

(HealthDay)—There is considerable variability associated with total knee arthroplasty (TKA) recommendations, according to a report published in the May issue of *Pain Medicine News*.

Liana Fraenkel, M.D., M.P.H., from the Yale School of Medicine in New Haven, Conn., and colleagues surveyed 406 <u>orthopedic surgeons</u> and 494 rheumatologists to examine whether they recommended TKA in various scenarios involving a 62-year-old patient with <u>knee osteoarthritis</u> and <u>moderate pain</u>.

The researchers found that for a patient described as having mild radiographic osteoarthritis and moderate pain, the proportion of



physicians who recommended TKA ranged from 30 to 55 percent. For a patient with moderate radiographic osteoarthritis and moderate pain, the proportion recommending TKA ranged from 39 to 71 percent. The proportion who recommended TKA varied significantly according to gender (59 percent for a male patient compared with 44 percent for a female patient). Compared with European surgeons (34 percent), U.S. surgeons were more likely to recommend TKA (52 percent). Rheumatologists and surgeons aged 40 years or younger were significantly more likely to recommend TKA (61 versus 46 percent for rheumatologists; 60 versus 48 percent for surgeons). Rheumatologists who were academics were significantly more likely to recommend TKA (63 versus 48 percent). Rheumatologists were also significantly more likely to recommend TKA (63 versus 48 percent). Rheumatologists were also significantly more likely to recommend TKA (63 versus 48 percent). Rheumatologists were also significantly more likely to recommend TKA (63 versus 48 percent). Rheumatologists were also significantly more likely to recommend TKA (63 versus 48 percent). Rheumatologists were also significantly more likely to recommend TKA (64 versus 48 percent). Rheumatologists were also significantly more likely to recommend TKA (64 versus 48 percent). Rheumatologists were also significantly more likely to recommend TKA for patients with moderate versus mild X-ray changes (60 versus 41 percent), and to retirees versus working people (56 versus 42 percent).

"Studies are needed to determine <u>appropriateness criteria</u> for TKA in order to reduce the unwarranted variability associated with TKA," Fraenkel said in a statement.

## More information: More Information

Health News Copyright © 2013 HealthDay. All rights reserved.

Citation: Marked variability seen in total knee arthroplasty indications (2013, June 14) retrieved 20 November 2023 from <u>https://medicalxpress.com/news/2013-06-variability-total-knee-arthroplasty-indications.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.