

Researchers discover how inhibitory neurons behave during critical periods of learning

August 25 2013

We've all heard the saying "you can't teach an old dog new tricks." Now neuroscientists are beginning to explain the science behind the adage.

For years, neuroscientists have struggled to understand how the microcircuitry of the brain makes learning easier for the young, and more difficult for the old. New findings published in the journal *Nature* by Carnegie Mellon University, the University of California, Los Angeles and the University of California, Irvine show how one component of the brain's [circuitry](#)—inhibitory neurons—behave during critical periods of learning. The paper is available online as an Advance Online Publication.

The brain is made up of two types of cells—inhibitory and excitatory neurons. Networks of these two kinds of neurons are responsible for processing sensory information like images, sounds and smells, and for [cognitive functioning](#). About 80 percent of neurons are excitatory. Traditional [scientific tools](#) only allowed scientists to study the excitatory neurons.

"We knew from previous studies that excitatory cells propagate information. We also knew that inhibitory neurons played a critical role in setting up heightened plasticity in the young, but ideas about what exactly those cells were doing were controversial. Since we couldn't study the cells, we could only hypothesize how they were behaving during critical learning periods," said Sandra J. Kuhlman, assistant professor of [biological sciences](#) at Carnegie Mellon and member of the

joint Carnegie Mellon/University of Pittsburgh Center for the Neural Basis of Cognition.

The prevailing theory on inhibitory neurons was that, as they mature, they reach an increased level of activity that fosters optimal periods of learning. But as the brain ages into adulthood and the inhibitory neurons continue to mature, they become even stronger to the point where they impede learning.

Newly developed genetic and [imaging technologies](#) are now allowing researchers to visualize inhibitory neurons in the brain and record their activity in response to a variety of stimuli. As a postdoctoral student at UCLA in the laboratory of Associate Professor of Neurobiology Joshua T. Trachtenberg, Kuhlman and her colleagues used these new techniques to record the activity of inhibitory neurons during critical learning periods. They found that, during heightened periods of learning, the [inhibitory neurons](#) didn't fire more as had been expected. They fired much less frequently—up to half as often.

"When you're young you haven't experienced much, so your brain needs to be a sponge that soaks up all types of information. It seems that the brain turns off the inhibitory cells in order to allow this to happen," Kuhlman said. "As adults we've already learned a great number of things, so our brains don't necessarily need to soak up every piece of information. This doesn't mean that adults can't learn, it just means when they learn, their neurons need to behave differently."

More information: Paper: <http://dx.doi.org/10.1038/nature12485>

Provided by Carnegie Mellon University

Citation: Researchers discover how inhibitory neurons behave during critical periods of learning (2013, August 25) retrieved 2 February 2024 from <https://medicalxpress.com/news/2013-08-inhibitory-neurons-critical-periods.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.