

# Pre-term infants with severe retinopathy more likely to have non-visual disabilities

February 4 2014

---

In a group of very low-birth-weight infants, severe retinopathy of prematurity was associated with nonvisual disabilities at age 5 years, according to a study in the February 5 issue of *JAMA*.

Severe retinopathy (disease of the retina) of prematurity occurs in [premature infants](#) treated with excessive concentrations of oxygen and is a serious complication of [neonatal intensive care](#) for [preterm infants](#). "Although the incidence of severe retinopathy has increased since the late 1980s, blindness caused by retinopathy has become rare in developed countries. Consequently, clinicians and parents may conclude that severe retinopathy is no longer associated with childhood impairments," according to background information in the article.

Barbara Schmidt, M.D., M.Sc., of Children's Hospital of Philadelphia, and colleagues investigated whether infants with severe retinopathy retain an increased risk of nonvisual disabilities compared with those without severe retinopathy. This analysis (using data from a trial, Caffeine for Apnea of Prematurity), included infants with birth weights between 1.1 and 2.8 lbs. who were born between 1999 and 2004 and followed-up at age 5 years (2005-2011).

Of 1,815 eligible infants, 1,582 (87 percent) had complete (n = 1,523) or partial (n = 59) 5-year assessments. Of 95 with severe retinopathy, 40 percent had at least 1 nonvisual disability at 5 years compared with 16 percent of children without it. Fourteen of 94 children (15 percent) with and 36 of 1,487 children (2.4 percent) without severe retinopathy had

more than 1 nonvisual disability. Motor impairment, cognitive impairment, and severe hearing loss were 3 to 4 times more common in children with severe retinopathy than those without severe retinopathy.

The authors write that these findings may help improve the ability to counsel parents and to select high-risk [infants](#) for long-term follow-up.

"Severe [retinopathy](#) of [prematurity](#) remains an adverse outcome of neonatal intensive care with poor prognosis for child development, although blindness can mostly be prevented by timely retinal therapy."

**More information:** [DOI: 10.1001/jama.282153](https://doi.org/10.1001/jama.282153)

Provided by The JAMA Network Journals

Citation: Pre-term infants with severe retinopathy more likely to have non-visual disabilities (2014, February 4) retrieved 27 February 2023 from <https://medicalxpress.com/news/2014-02-pre-term-infants-severe-retinopathy-non-visual.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--