

Finding keys to glioblastoma therapeutic resistance

August 26 2014

Researchers at the University of California, San Diego School of Medicine have found one of the keys to why certain glioblastomas – the primary form of a deadly brain cancer – are resistant to drug therapy. The answer lies not in the DNA sequence of the tumor, but in its epigenetic signature. These findings have been published online as a priority report in the journal *Oncotarget*.

"There is a growing interest to guide cancer therapy by sequencing the DNA of the cancer cell," said Clark Chen, MD, PhD, vice-chairman of Research and Academic Development, UC San Diego Division of Neurosurgery and the principal investigator of the study. "Our study demonstrates that the sensitivity of glioblastoma to a drug is influenced not only by the content of its DNA sequences, but also by how the DNA sequences are organized and interpreted by the cell."

The team of scientists, led by Chen, used a method called comparative gene signature analysis to study the genetic profiles of tumor specimens collected from approximately 900 glioblastoma patients. The method allows investigators to discriminate whether specific [cellular processes](#) are "turned on" or "turned off" in [glioblastomas](#). "Our study showed that not all glioblastomas are the same. We were able to classify glioblastomas based on the type of cellular processes that the [cancer cells](#) used to drive tumor growth," said Jie Li, PhD, senior postdoctoral researcher in the Center for Theoretical and Applied Neuro-Oncology at UC San Diego and co-first author of the paper.

One of these cellular processes involves Epidermal Growth Factor Receptor (EGFR). The study revealed that EGFR signaling is suppressed in a subset of glioblastomas. Importantly, this suppression is not the result of altered DNA sequences or mutations. Instead, EGFR is turned off as a result of how the DNA encoding the EGFR gene is organized in the cancer cell. This form of regulation is termed "epigenetic." Because EGFR is turned off in these glioblastomas, they become insensitive to drugs designed to inhibit EGFR signaling.

"Our research suggests that the selection of appropriate therapies for our brain [tumor](#) patients will require a meaningful synthesis of genetic and epigenetic information derived from the cancer cell," said co-first author Zachary J. Taich.

Provided by University of California - San Diego

Citation: Finding keys to glioblastoma therapeutic resistance (2014, August 26) retrieved 3 February 2024 from <https://medicalxpress.com/news/2014-08-keys-glioblastoma-therapeutic-resistance.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--