

# Research shows 25 percent fewer opioid-related deaths in states allowing medical marijuana

August 25 2014

---

On average, states allowing the medical use of marijuana have lower rates of deaths resulting from opioid analgesic overdoses than states without such laws. Opioid analgesics, such as OxyContin, Percocet and Vicodin, are prescribed for moderate to severe pain, and work by suppressing a person's perception of pain.

A new multi-institutional study, published in *JAMA Internal Medicine* and led by researchers at the Perelman School of Medicine at the University of Pennsylvania, examined the rate of deaths caused by [opioid](#) overdoses between 1999 and 2010. Results reveal that on average, the 13 [states](#) allowing the use of [medical marijuana](#) had a 24.8 percent lower annual opioid overdose mortality rate after the laws were enacted than states without the laws, indicating that the alternative treatment may be safer for patients suffering from [chronic pain](#) related to cancer and other conditions.

Approximately 60 percent of all deaths resulting from opioid analgesic overdoses occur in patients who have legitimate prescriptions. Additionally, the proportion of patients in the United States who are prescribed opioids for non-cancer pain has almost doubled over the past decade, indicating the need to do a more focused examination on the safety and efficacy of these and other treatment options. In states allowing the use of medical cannabis, the drugs may be prescribed as an alternative to opioids.

While noting that evidence for the pain-relieving properties of cannabis is limited, some studies have suggested "it may provide relief for some individuals," said lead author, Marcus A. Bachhuber, MD, Robert Wood Johnson Foundation Clinical Scholar at Penn and the Philadelphia VA Medical Center. "In addition, people already taking opioids for pain may supplement with medical marijuana and be able to lower their painkiller dose, thus lowering their risk of overdose.

Additional results of the study show that the relationship between lower opioid [overdose deaths](#) and medical marijuana laws strengthened over time; deaths were nearly 20 percent lower in the first year after a state's law was implemented, and 33.7 percent lower five years after implementation. While safer treatment of chronic pain may help to explain lower rates of overdose deaths, medical marijuana laws may also change the way people misuse or abuse opioid painkillers, as marijuana and opioids stimulate similar areas in the brain's pathways. The authors suggest that as more states implement medical [marijuana laws](#), future studies should examine the association between such laws and opioid overdoses to confirm their findings.

**More information:** *JAMA Intern Med.* Published online August 25, 2014. [DOI: 10.1001/jamainternmed.2014.4005](https://doi.org/10.1001/jamainternmed.2014.4005)  
*JAMA Intern Med.* Published online August 18, 2014. [DOI: 10.1001/jamainternmed.2014.2716](https://doi.org/10.1001/jamainternmed.2014.2716)

Provided by University of Pennsylvania School of Medicine

Citation: Research shows 25 percent fewer opioid-related deaths in states allowing medical marijuana (2014, August 25) retrieved 3 February 2024 from <https://medicalxpress.com/news/2014-08-percent-opioid-related-deaths-states-medical.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.