


Blood test for 'nicked' protein predicts prostate cancer treatment response

September 3 2014



Micrograph showing prostatic acinar adenocarcinoma (the most common form of prostate cancer) Credit: Wikipedia, [CC BY-SA 3.0](https://creativecommons.org/licenses/by-sa/3.0/)

Prostate cancer patients whose tumors contain a shortened protein called AR-V7, which can be detected in the blood, are less likely to respond to two widely used drugs for metastatic prostate cancer, according to results of a study led by researchers at the Johns Hopkins Kimmel Cancer Center. If large-scale studies validate the findings, the investigators say men with detectable blood levels of AR-V7 should avoid these two drugs

and instead take other medicines to treat their prostate cancer. A report on the work is described online Sept. 3 in the *New England Journal of Medicine*.

The study evaluated two groups of 31 men with [prostate cancer](#) that had spread and whose blood levels of prostate-specific antigen (PSA) were still rising despite low testosterone levels. Investigators gave each man either enzalutamide (Xtandi) or abiraterone (Zytiga) and tracked whether their PSA levels continued to rise, an indication that the drugs were not working. In the enzalutamide group, none of 12 patients whose [blood samples](#) tested positive for AR-V7 responded to the drug, compared with 10 responders among 19 men who had no AR-V7 detected. In the abiraterone group, none of six AR-V7-positive patients responded, compared with 17 responders among 25 patients lacking AR-V7.

Enzalutamide and abiraterone have been very successful in lengthening the lives of about 80 percent of patients with [metastatic prostate cancer](#), says Emmanuel Antonarakis, M.D., assistant professor of oncology at Johns Hopkins, but the drugs do not work in the remaining 20 percent of patients.

"Until now, we haven't been able to predict which patients will not respond to these therapies. If our results are confirmed by other researchers, a blood test could use AR-V7 as a biomarker to predict enzalutamide and abiraterone resistance, and let us direct patients who test positive for AR-V7 toward other types of therapy sooner, saving time and money while avoiding futile therapy," says Antonarakis.

Prostate cancer thrives on male sex hormones (or "androgens"), including testosterone. Enzalutamide and abiraterone target proteins called androgen receptors and block the receptors' ability to activate prostate [cancer cells](#). AR-V7 is a shortened form of the androgen receptor that lacks a binding spot targeted by enzalutamide and

abiraterone. With no binding spot for the two drugs, AR-V7 is free to manipulate [prostate cancer cells'](#) genetic material, which makes the cancer cells grow and spread.

Antonarakis and his colleague Jun Luo, Ph.D., who first identified AR-V7 in 2008, also tracked patients' progression-free survival (the length of time a patient lives with the disease but does not get worse) and overall survival. They found that, in men receiving enzalutamide, progression-free survival was 2.1 months in AR-V7-positive patients and 6.1 months in AR-V7-negative patients, while overall survival was 5.5 months in AR-V7-positive men and up to 9 months in AR-V7-negative men. Similarly, in men receiving abiraterone, progression-free survival was 2.3 months in AR-V7-positive patients and up to 6 months in AR-V7-negative patients, while overall survival was 10.6 months in AR-V7-positive men and up to 12 months in AR-V7-negative men. The investigators caution that most of the study patients had advanced disease and received multiple prior therapies, so their outcomes may not be generalizable to all [men](#) with prostate cancer.

"Patients whose blood samples contained AR-V7 got no benefit from either enzalutamide or abiraterone," says Antonarakis. He adds that the shortened AR-V7 protein could appear in patients' blood samples at the very start of therapy or acquired later, after therapy has begun. He says, "This test could be used before starting enzalutamide or abiraterone therapy, and if the test shows the presence of AR-V7, patients may opt for a different therapy. It could also be used to monitor [patients](#) receiving [enzalutamide](#) or [abiraterone](#) for AR-V7, providing an indication these drugs may not work for much longer."

Provided by Johns Hopkins University School of Medicine

Citation: Blood test for 'nicked' protein predicts prostate cancer treatment response (2014,

September 3) retrieved 3 February 2024 from <https://medicalxpress.com/news/2014-09-blood-nicked-protein-prostate-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.