

Another breastfeeding benefit: Preparing baby's belly for solid food

February 5 2015

The moment of birth marks the beginning of a beautiful, lifelong relationship between a baby and the billions of microbes that will soon colonize his or her gastrointestinal tract.

In a study published today in *Frontiers in Cellular and Infection Microbiology*, researchers from the UNC School of Medicine and UNC College of Arts and Sciences found that a baby's diet during the first few months of life has a profound influence on the composition, diversity, and stability of the gut microbiome. These factors, in turn, influence the baby's ability to transition from milk to solid foods and may have long-term health effects.

"We found that babies who are fed only [breast milk](#) have [microbial communities](#) that seem more ready for the introduction of solid foods," said Andrea Azcarate-Peril, PhD, assistant professor in the department of cell biology and physiology and the study's senior author. "The transition to solids is much more dramatic for the microbiomes of babies that are not exclusively breastfed. We think the microbiomes of non-exclusively [breastfed babies](#) could contribute to more stomach aches and colic."

The discovery adds to the growing awareness that the gut microbiome plays a major role in helping us digest food and fight pathogens, among other functions.

"This study provides yet more support for recommendations by the

World Health Organization and others to breastfeed exclusively during the first six months of life," said Amanda Thompson, PhD, associate professor in the department of anthropology, a Carolina Population Center faculty fellow, and the study's first author. "We can see from the data that including formula in an infant's diet does change the gut bacteria even if you are also breastfeeding. Exclusive breastfeeding seems to really smooth out the transition to solid foods."

For this study, the research team collected [stool samples](#) and information about the diets and health of nine babies as they grew from ages 2 weeks to 14 months. Applying genomic sequencing techniques to the stool samples, the scientists deduced the types and functions of the bacteria in the babies' gut microbiomes.

The analysis revealed that during the first few months of life there were clear differences between the microbiomes of babies that were exclusively breastfed as compared to those fed both formula and breast milk. This finding is consistent with previous studies.

What surprised Thompson and Azcarate-Peril, who is the director of the UNC Microbiome Core Facility, was the drastic genetic differences in stool samples taken after babies began eating solid food.

Researchers found differing amounts of about 20 bacterial enzymes in exclusively breastfed babies when compared to exclusively breastfed babies that received solid food. This indicated that some new bacterial species had entered the scene to help process the new food types. In babies fed both formula and breast milk - and then introduced to solid foods - the samples revealed about 230 enzymes, indicating a much more dramatic shift in microbial composition.

The microbiomes of exclusively breastfed babies tended to be less diverse and were dominated by Bifidobacterium, a type of bacteria

considered beneficial for digestion. Babies fed a mixture of breast milk and formula had a lower proportion of Bifidobacterium.

The study suggests that the makeup of the microbiome can affect a baby's ability to digest food in the short term and potentially influence long-term health.

Although microbiome research is still in its early stages, gut microbes are thought to potentially play a role in obesity, allergies, and gastrointestinal problems, such as irritable bowel syndrome.

"The study advances our understanding of how the [gut microbiome](#) develops early in life, which is clearly a really important time period for a person's current and future health," said Thompson.

The researchers also compared the microbiomes of [babies](#) that attended daycare to those that stayed in the home. Attending daycare was also associated with more diverse microbial communities overall, but feeding practices remained the most important factor influencing how the microbiome responded to the introduction of solid foods.

Provided by University of North Carolina Health Care

Citation: Another breastfeeding benefit: Preparing baby's belly for solid food (2015, February 5) retrieved 20 July 2023 from <https://medicalxpress.com/news/2015-02-breastfeeding-benefit-baby-belly-solid.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--