

# More than one-third of perimenopausal women develop insomnia

October 14 2015

---

Millions of women may likely be sleep-deprived. It's already a known fact that women are more predisposed to insomnia. Now a new study presented at the 2015 Annual Meeting of The North American Menopause Society (NAMS) earlier this month suggests that perimenopausal women have an even greater risk for developing insomnia. Considering that perimenopause will affect roughly 500 million women within the next decade, that's a lot of tired women.

What's worse is that the study found that [insomnia](#) symptoms are likely to get worse and more prevalent in the later stages of perimenopause (the transition period to menopause). In fact, the odds of having any one symptom of insomnia were 1.3 times greater for those in late stage versus early stage of perimenopause. The odds of developing [chronic insomnia](#) were 1.5 times greater for those in perimenopause than [pre-perimenopause](#).

"We found that there was a lot of research regarding insomnia in general but very little that addressed the insomnia trajectory in one of the higher risk groups of women—those transitioning to menopause," says Dr. Colleen Ciano of the College of Nursing at The Pennsylvania State University and lead author of the study.

The most common sleep-related complaints include: difficulty falling asleep, waking after sleep onset, and sleep quality. Of the 3,302 study participants, more than one-third suffered from insomnia, reporting "awakenings" as the most frequent insomnia symptom.

"Given the strong link between insomnia and such poor health outcomes as heart disease and obesity, this study offers valuable insight for physicians who are treating middle-aged patients and considering various preventative treatment options," says Dr. Wulf Utian, NAMS executive director.

Provided by The North American Menopause Society

Citation: More than one-third of perimenopausal women develop insomnia (2015, October 14) retrieved 29 March 2023 from <https://medicalxpress.com/news/2015-10-one-third-perimenopausal-women-insomnia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.