

Durable response with insulin pump therapy in T2DM

February 12 2016



(HealthDay)—Insulin pump therapy is more effective than multiple

daily injections (MDI) for glycemic control in patients with type 2 diabetes, according to a study published online Feb. 8 in *Diabetes, Obesity and Metabolism*.

Ronnie Aronson, M.D., from LMC Diabetes & Endocrinology in Toronto, and colleagues compared insulin pump therapy and MDI in 331 [patients](#) with type 2 diabetes. Participants with glycated [hemoglobin](#) ≥ 8.0 percent and ≤ 12 percent were randomly allocated to pump therapy or continued MDI in a six-month randomization phase (RP). During a six-month continuation phase (CP), the MDI group was switched to pump therapy.

The researchers found that the reduction in glycated hemoglobin was significantly greater with pump therapy versus MDI at the end of the RP (-1.1 ± 1.2 versus -0.4 ± 1.1 percent; P

"Patients with refractory hyperglycemia on a current basal-prandial injection regimen should be considered appropriate candidates for pump therapy, and may obtain sustained [glycemic control](#) with a favorable safety profile and reduction of insulin dose," the authors write.

Several authors disclosed financial ties to pharmaceutical and medical technology companies, including Medtronic, which funded the study; several authors disclosed full-time employment by Medtronic.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Durable response with insulin pump therapy in T2DM (2016, February 12) retrieved 2 February 2024 from <https://medicalxpress.com/news/2016-02-durable-response-insulin-therapy->

[t2dm.html](#)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.