

Testosterone undecanoate improves sexual function in men with type 2 diabetes

May 6 2016

In a recent placebo-controlled study, long acting testosterone undecanoate (an ester of testosterone) improved erectile function, intercourse satisfaction, and sexual desire scores in type 2 diabetic men with severe hypogonadism, a condition in which the body doesn't produce enough testosterone.

Only sexual desire improved significantly with [testosterone](#) replacement therapy in those with mild hypogonadism.

"The study's results also suggest that trials of [testosterone therapy](#) should be for a minimum of 6 months and not shorter periods as suggested by some guidelines," said Dr. Geoffrey Hackett, lead author of the *BJU International* study.

More information: Geoffrey Hackett et al, Testosterone Undecanoate improves Sexual Function in Men with Type 2 diabetes and Severe Hypogonadism: Results from a 30 week randomized placebo controlled study, *BJU International* (2016). [DOI: 10.1111/bju.13516](https://doi.org/10.1111/bju.13516)

Provided by Wiley

Citation: Testosterone undecanoate improves sexual function in men with type 2 diabetes (2016, May 6) retrieved 18 January 2024 from <https://medicalxpress.com/news/2016-05-testosterone-undecanoate-sexual-function-men.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.