


# New discovery toward developing Huntington's disease treatments before symptoms appear

February 22 2017

---



A montage of three images of single striatal neurons transfected with a disease-associated version of huntingtin, the protein that causes Huntington's disease. Nuclei of untransfected neurons are seen in the background (blue). The neuron in the center (yellow) contains an abnormal intracellular accumulation of

huntingtin called an inclusion body (orange). Credit: Wikipedia/ Creative Commons Attribution 3.0 Unported license

Early warning signs of Huntington's disease have been uncovered in a sheep carrying the human HD mutation, leading the way for new insight into this devastating illness, a new study in *Scientific Reports* has found.

Researchers from the University of Surrey and University of Cambridge have identified early biomarkers of Huntington's [disease](#) during examinations of HD [sheep](#) still at a pre-symptomatic stage of the disease.

Up until this point, the five-year-old HD sheep displayed no signs of the illness, but this comprehensive study identified clear metabolic changes in the affected animals. These new findings reveal that Huntington's disease affects important metabolic processes in the body prior to the appearance of [physical symptoms](#).

Huntington's disease is a genetic neurodegenerative disorder caused by the expansion of a CAG repeat in the HD gene. The disease affects more than 6,700 people in the UK. Huntington's disease is incurable, and patients typically die 10-25 years after diagnosis. The sheep model of Huntington's disease has been developed to increase knowledge about Huntington's disease.

During this study, [blood samples](#) were taken from normal and HD animals every two hours over a 24-hour period and their metabolic profiles assessed using a targeted metabolomics approach established at the University of Surrey. Unlike previous research in this area, which was affected by external environmental factors that impacted upon metabolic profiling, sheep in this study were monitored in a well-

controlled setting, negating any outside influences.

Blood measurements found startling differences in the biochemistry of the sheep carrying the HD gene, compared to the normal sheep. Significant changes were observed in 89 of the 130 metabolites measured in their blood, with increased levels of the amino acids, arginine and citrulline, and decreases in sphingolipids and fatty acids that are commonly found in brain and nervous tissue.

The alterations in these metabolites, which include key components of the urea cycle and nitric oxide pathways (both vital body processes), suggest that both of these processes are dysregulated in the early stages of Huntington's disease, and that the illness affects the body long before physical symptoms appear.

The identification of these biomarkers may help to track disease in pre-symptomatic patients, and could help researchers develop strategies to remedy the biochemical abnormalities.

Professor Debra Skene from the University of Surrey said: "Metabolic profiling has revealed novel biomarkers that will be useful to monitor Huntington's disease progression.

"Our research shows that this disease affects the body in a number of ways before the tell-tale signs of Huntington's disease become visible. This is not the end of our research, we will continue to observe these sheep and study how this disease evolves."

Professor Jenny Morton from the University of Cambridge said: "Despite its devastating impacts on patients and their families, there are currently limited treatments options, and no cure for Huntington's disease.

"The development of objective and reliable biomarkers that can be rapidly measured from blood samples becomes immeasurably important once clinical trials for therapies begin.

"The more we learn about this devastating illness the better chance we have of finding a cure."

**More information:** Debra J. Skene et al. Metabolic profiling of presymptomatic Huntington's disease sheep reveals novel biomarkers, *Scientific Reports* (2017). [DOI: 10.1038/srep43030](https://doi.org/10.1038/srep43030)

Provided by University of Surrey

Citation: New discovery toward developing Huntington's disease treatments before symptoms appear (2017, February 22) retrieved 20 September 2024 from <https://medicalxpress.com/news/2017-02-discovery-huntington-disease-treatments-symptoms.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.