

Neuronal stimulation regulates appetite and glucose levels in mice

February 21 2017

Drugs that target signaling by glucagon-like peptide (GLP-1) in the gut and pancreas are commonly used to regulate hypoglycemia in patients with type II diabetes. However, these drugs are associated with a number of unpleasant side effects that may be linked to other targets of GLP-1 signaling. A subset of neurons in the brain also produces GLP-1 and related peptides, but how brain-mediated GLP-1 signaling influences metabolism and appetite is not clear.

This week in the *JCI*, a study led by Michael Scott at the University of Virginia explores how stimulation of GLP-1-producing neurons can control appetite and glucose levels in mice.

The researchers selectively excited GLP-1 neurons in the hypothalamus of mice and found that the stimulation reduced animals' [food intake](#) and suppressed the generation of glucose from energy reserves.

However, the stimulation's effects on glucose handling varied between lean and obese mice—in obese mice, exciting GLP-1 neurons suppressed [appetite](#) without changing [glucose levels](#).

The finding suggests that distinct pathways with different sensitivities to individual metabolic state may mediate the behavioral and physiological effects of GLP-1 signaling.

More information: Ronald P. Gaykema et al, Activation of murine pre-proglucagon-producing neurons reduces food intake and body

weight, *Journal of Clinical Investigation* (2017). [DOI: 10.1172/JCI81335](https://doi.org/10.1172/JCI81335)

Provided by JCI Journals

Citation: Neuronal stimulation regulates appetite and glucose levels in mice (2017, February 21) retrieved 3 July 2023 from <https://medicalxpress.com/news/2017-02-neuronal-appetite-glucose-mice.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.