

Nivolumab produces durable responses and long-term survival in severe liver cancer patients

April 21 2017

Results from the CheckMate 040 study presented today found that nivolumab, an immuno-oncology drug which acts by modulating the immune system, produces durable responses with long-term survival rates, regardless of whether or not patients were infected with Hepatitis B or C. Interim results from the study, presented at The International Liver Congress 2017 in Amsterdam, The Netherlands, showed that the overall objective response rate (ORR) by blinded independent central review (BICR) was 14.5% and ORR by investigator assessment was 19.3% in sorafenib-experienced patients in the dose expansion phase of CheckMate 040. Responses by BICR were ongoing in 71.4% (15/21) of patients, and the 12-month overall survival rate in this cohort was 59.9%. The safety profile of nivolumab was manageable and consistent with that reported in other tumour types.

Liver cancer, or hepatocellular carcinoma (HCC), is the second most common cause of cancer-related deaths worldwide.^{1,2} The prognosis for patients with advanced [liver](#) cancer is poor,² and the multikinase inhibitor, sorafenib, is the only approved systemic treatment.³ If the patient is not tolerant or has contraindications for sorafenib therapy, there is currently no standard of care and therefore patients lack effective treatment options.³ Nivolumab has already increased survival time in different types of cancers, and has become an important treatment option for certain types of kidney, blood, melanoma and non-small cell lung cancer.⁴ Preliminary results from the CheckMate 040

study presented earlier this year suggested that [nivolumab](#) could be an option for the treatment of liver cancer.⁵ Nivolumab is not yet licensed for HCC in the EU.

"The durable responses and survival rates that were achieved with nivolumab are very welcome, especially as the side effects were manageable," said Prof Bruno Sangro, Head of Liver Unit, Clínica Universidad de Navarra and CIBEREHD, and study author. "These data support the potential of nivolumab in the treatment and stabilisation of advanced liver cancer in those patients who have progressed on sorafenib, with or without chronic viral hepatitis."

The CheckMate 040 study is a Phase 1/2, multi-cohort, open-label study of nivolumab conducted in patients with advanced liver cancer who were not suitable for surgery.⁶ The primary endpoint of the study was ORR by blinded, independent central review. All 145 patients previously treated with sorafenib in the dose-expansion portion of the study were given intravenous nivolumab 3 mg/kg every 2 weeks until the cancer progressed or side effects became intolerable.

Of the 145 patients who had previously received sorafenib, 132 (91.0%) had progression of their cancer and 12 (8.3%) were intolerant of the therapy. The median follow up was 12.9 months in this interim analysis of the dose expansion phase. The median duration of response (DOR) was not yet reached, and 8/21 responders had a DOR of greater than 12 months. The overall median overall survival (OS) rate was 16.7 months, and it was not reached in those with chronic viral hepatitis B and C. Responses to nivolumab occurred regardless of programmed death-1 (PD-1) ligand expression on tumour cells. Overall, grade 3/4 treatment-related adverse events occurred in 16.6% of patients.

Nivolumab is a programmed-death-1 (PD-1) immune checkpoint inhibitor that is designed to use the body's own immune system to help

restore the anti-cancer immune response.⁵ It restores T-cell-mediated anti-tumour activity so that the T cells recognise and attack [cancer](#) cells.⁵

"The reported median survival of 16.7 months in patients previously treated with [sorafenib](#) is promising and it encourages the evaluation of nivolumab in [patients](#) affected with [hepatocellular carcinoma](#)," said Prof Alejandro Forner, BCLC group, Liver Unit, Hospital Clinic Barcelona, Spain and member of the EASL Governing Board.

More information: Abstract: Nivolumab in sorafenib-experienced patients with advanced hepatocellular carcinoma (HCC) with or without chronic viral hepatitis: CheckMate 040 study (GS010), The International Liver Congress 2017.

References:

- 1 World Health Organization. Cancer. Available from: www.who.int/mediacentre/factsheets/fs297/en/. Last accessed: April 2017.
- 2 World Health Organization. GLOBOCAN 2012: Estimated cancer incidence, mortality and prevalence worldwide in 2012. Available from: globocan.iarc.fr/Pages/fact_sheets_cancer.aspx. Last accessed: April 2017.
- 3 EASL-EORTC Clinical Practice Guidelines: Management of hepatocellular carcinoma. J Hepatol. 2012;56:908-943.
- 4 European Medicines Agency. EPAR summary for the public: Opvido (nivolumab). Available from: www.ema.europa.eu/docs/en_GB/d...3985/WC500189768.pdf. Last accessed: April 2017.

5 Melero I, et al. Nivolumab dose escalation and expansion in patients with advanced hepatocellular carcinoma (HCC): The CheckMate 040 study. J Clin Oncol. 2017;35 (suppl 4S: abstract 226.

6 ClinicalTrials.gov. NCT Identifier: NCT01658878. A study to evaluate the effectiveness, safety and tolerability of nivolumab and the combination nivolumab plus ipilimumab in patients with advanced liver cancer (CheckMate 040). Available from: clinicaltrials.gov/show/NCT01658878. Last accessed: April 2017.

7 Bristol-Myers Squibb. How does OPDIVO work inside my body? Available from: [www.opdivo.com/advanced-renal- ... ivo/how-opdivo-works](http://www.opdivo.com/advanced-renal-...ivo/how-opdivo-works). Last accessed: April 2017.

Provided by European Association for the Study of the Liver

Citation: Nivolumab produces durable responses and long-term survival in severe liver cancer patients (2017, April 21) retrieved 4 February 2024 from <https://medicalxpress.com/news/2017-04-nivolumab-durable-responses-long-term-survival.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.