

A potential cure for metastatic prostate cancer? Treatment combination shows early promise

April 18 2017

In the past, all forms of metastatic prostate cancer have been considered incurable. In recent years, the FDA has approved six drugs for men with metastatic disease, all of which can increase survival. In a study published in *Urology*, researchers demonstrate for the first time that an aggressive combination of systemic therapy (drug treatment) with local therapy (surgery and radiation) directed at both the primary tumor and metastasis can eliminate all detectable disease in selected patients with metastatic prostate cancer.

While the study is only a first step, one-fifth of the [patients](#) treated had no detectable disease, with an undetectable prostate-specific-antigen (PSA) and normal blood testosterone, after 20 months. The results suggest that some men who have previously been considered incurable can possibly be cured; investigators also establish a new paradigm for testing various drug combinations in conjunction with local treatment of the prostate to determine which is the best approach (ie, has the highest undetectable disease rate). Such results could not have been achieved with any single [therapy](#) alone.

According to lead investigator Howard I. Scher, MD, Chief of the Genitourinary Oncology Service at Memorial Sloan Kettering Cancer Center in New York City, "The sequential use of the three different modalities helped illustrate the role and importance of each in achieving the undetectable PSA with normal testosterone level end point, which

represents a 'no-evidence of disease' status." Longer follow-up is needed to determine whether these patients were in fact cured.

Twenty men with metastatic [prostate cancer](#), five with extra-pelvic lymph nodal disease and 15 with bone with or without nodal disease, were treated with androgen deprivation therapy (ADT), radical [surgery](#) that included a retroperitoneal [lymph node dissection](#) as needed, and radiation therapy to visible metastatic lesions in bone. ADT was stopped after a minimum of six months if an undetectable PSA was achieved after combined modality therapy. Other patients were treated continuously.

The combined treatment regimen including surgery was well tolerated. Matthew J. O'Shaughnessy, MD, PhD, Urology Service, Department of Surgery, Memorial Sloan Kettering Cancer Center, commented "While the role of local therapy in [metastatic prostate cancer](#) is still under investigation, aggressive resection of visible disease performed by experienced surgeons was critical to the outcome."

Of the five patients with extra-pelvic lymph node involvement, four achieved an undetectable PSA after ADT and surgery, while the fifth needed radiation to reach this milestone. However, none achieved the primary end point of undetectable PSA with testosterone recovery at 20 months after initiation of therapy with ADT alone, although one patient had a PSA of

Citation: A potential cure for metastatic prostate cancer? Treatment combination shows early promise (2017, April 18) retrieved 23 July 2023 from <https://medicalxpress.com/news/2017-04-potential-metastatic-prostate-cancer-treatment.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.