

Making prosthetic limbs feel more natural

May 31 2017



MIT researchers are developing a new surgical approach that would allow amputees to receive sensory feedback from their prosthetic limbs and improve their ability to control them. Credit: Jose-Luis Olivares

A new surgical technique devised by MIT researchers could allow prosthetic limbs to feel much more like natural limbs. Through coordination of the patient's prosthetic limb, existing nerves, and muscle

grafts, amputees would be able to sense where their limbs are in space and to feel how much force is being applied to them.

This type of system could help to reduce the rejection rate of prosthetic limbs, which is around 20 percent.

"We're talking about a dramatic improvement in patient care," says Hugh Herr, a professor of media arts and sciences and the senior author of the study. "Right now there's no robust neural method for a person with [limb amputation](#) to feel proprioceptive positions and forces applied to the prosthesis. Imagine how that would completely hinder one's ability to move, to successfully balance, or to manipulate objects."

In the new study, which appears in *Science Robotics* on May 31, the researchers demonstrated in rats that their technique generates [muscle](#)-tendon sensory feedback to the nervous system, which should be able to convey information about a [prosthetic limb](#)'s placement and the forces applied to it. They now plan to begin implementing this approach in human amputees, including Herr, whose legs were amputated below the knee when he was 17.

Shriya Srinivasan, a graduate student in the Harvard-MIT Program in Health Sciences and Technology (HST), is the paper's lead author. Other authors are Media Lab visiting scientist Matthew Carty, MIT undergraduate Peter Calvaresi, HST graduate students Tyler Clites and Benjamin Maimon, Media Lab graduate student Cameron Taylor, and recent PhD recipient Anthony Zorzos.

Better feedback


During a conventional limb amputation, muscles are severed in a way that cuts off a key relationship that normally helps people control their limbs and sense where they are in space. Most muscles that control limb

movement occur in pairs known as agonist-antagonist pairs, such that one muscle stretches when the other contracts. For example, when you bend your elbow, the biceps muscle contracts, causing the triceps to stretch, and that triceps stretch sends sensory information related to position, velocity, and force back to the brain. The agonist-antagonist muscle relationship is also what allows people to independently control position and stiffness at their limb joints.

Without these intact muscle pairs, persons with limb amputation have no way of sensing where their artificial limbs are, nor can they sense the forces applied to those limbs.

"They have to visually follow their hands or their limbs, because there isn't any feedback from the device or residual limb that tells their brain where their prosthetic limbs are in space," Srinivasan says.

The MIT team set out to recreate these agonist-antagonist muscle relationships. In many amputees, the nerves that send signals to the amputated limb remain intact. The researchers decided to take advantage of those nerves by connecting them to muscle pairs grafted from another part of the body into the amputation site.



The agonist-antagonist myoplastic interface (AMI) is an advanced neural interfacing construct for amputation surgeries to enable robust bidirectional communication with prostheses and provide naturally-generated sensory feedback information to the user. Credit: Stephanie Ku


These grafts, which would be about 4 centimeters by 1.5 centimeters in humans, consist of a pair of muscles that work together just like natural muscles. When the brain sends signals instructing a limb to move, one of the grafted muscles will contract, and its agonist will extend. The agonist muscle then sends feedback to the brain about how much the muscle moved and the forces applied to it.

In the *Science Robotics* paper, the researchers tested the muscle grafts in rats and found that when the rats contracted one muscle of the pair, the other muscle would move in the opposite way and send sensory information back to the brain.

Control system

In other work, the researchers have developed the components of a control system that will translate the nerve signals into instructions for moving the prosthetic limb. When the brain sends nerve impulses to the regenerated muscles, those signals will also be received by a microprocessor that controls the movement of the [artificial limb](#).

Neural stimulations will cause the agonist muscle to contract and the antagonist muscle to stretch. The stretched muscle will then provide neural feedback to allow the patient to feel where his or her limb is in space. The researchers expect that the brain will be able to rapidly learn how much control it has to exert to make an artificial limb move in the desired way.



Muscle histology stains confirming nerve regeneration and development of major muscle structures in the graft. Credit: Srinivasan et al., *Sci. Robot.* 2, eaan2971

"Using this framework, the patient will not have to think about how to control their artificial limb. When a patient imagines moving their phantom [limb](#), signals will be sent through nerves to the surgically constructed muscle pairs. Implanted muscle electrodes will then sense these signals for the control of synthetic motors in the external prosthesis," Herr says. "We think that because the brain is so good at remapping and it's so plastic, it will quickly adapt to knowing how much it has to contract each muscle graft for natural prosthetic control."

This type of feedback system should also allow people with a prosthetic arm, for example, to feel a torque applied to the prosthesis. "If you were to give a prosthetic-arm user a barbell to hold, they would actually feel the torque on the prosthetic wrist joint," Herr says.

The researchers anticipate that this strategy could work for nearly any amputee, including people whose amputations were performed many years ago.

"For almost any amputation scenario, as long as we have a little bit of the healthy nerve left, we can take that and put it into regenerative muscle grafts. We can harvest these muscle grafts from almost anywhere in the body, making this applicable to a large number of cases ranging from trauma to chronic pain," Srinivasan says.

More information: S.S. Srinivasan et al., "On Prosthetic Control: A Regenerative Agonist-Antagonist Myoneural Interface," *Science*

Robotics (2017). robotics.sciencemag.org/lookup.../scirobotics.aan2971

Provided by Massachusetts Institute of Technology

Citation: Making prosthetic limbs feel more natural (2017, May 31) retrieved 1 February 2024 from <https://medicalxpress.com/news/2017-05-prosthetic-limbs-natural.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.