


How much does dementia risk increase after traumatic brain injury?

July 5 2017



The WHO has predicted that TBI will become a leading cause of death and long-term illness during the next ten years. Already one per cent of the population in the United States suffers from a long-term disability caused by TBI. Credit: R Raj (University of Helsinki)

Having a moderate or severe traumatic brain injury (TBI) in middle-age is associated with an increased risk for future dementia, according to a retrospective study of more than 40,000 TBI patients published in *PLOS Medicine* by Rahul Raj of the University of Helsinki, Finland, and colleagues.


A TBI usually results from a blow or jolt to the head and, in more severe cases, can lead to torn tissues, bleeding and other physical damage to the brain. In the new study, researchers identified everyone in Finland between the ages of 18 and 65 who was hospitalized with a mild or a moderate-to-severe TBI between 1986 and 2014. They then used the Finnish Care Register for Health Care to track hospitalization, after the TBI, with dementia, Parkinson's disease, and amyotrophic lateral sclerosis (ALS) for all 40,639 patients.

3.5% of patients who had a moderate-to-severe TBI went on to receive a diagnosis of dementia, compared to 1.6% for those who only had a mild TBI. After adjusting for confounding factors, people with a moderate-to-severe TBI were 90% more likely than those with mild TBI to be diagnosed with dementia (hazard ratio [HR] 1.9; 95% confidence interval [CI] 1.6-2.2). The risk appeared to be most increased among those whose injury occurred in middle age (HR 2.7 [CI 1.8-4.2] for age 41-50, and 2.0 [CI 1.6-2.4] for age 51-60). There was no increase in the risk of Parkinson's disease or ALS with TBI. The study was limited by a significant number of deaths in the group of patients with moderate-to-severe TBI, potential misdiagnoses of TBIs, and the fact that the database only included patients who were hospitalized, for TBIs and subsequently with dementia or other neurodegenerative conditions, so may have missed those treated in an outpatient setting.

"Our results imply that in working-aged persons, moderate-to-severe TBI seems to increase the risk for developing neurodegenerative disease later in life," the authors say. "The effect of covariates, such as comorbidities,

life-style factors and the effect of genetics, should be accounted for in future etiological studies, as well as to improve diagnostics and prevention of dementia after TBI."

**RISK OF DEMENTIA IN WORKING-AGE PEOPLE
AFTER MODERATE-TO-SEVERE TBI COMPARED TO MILD TBI**



The risk of a memory disorder was significantly higher among patients with medium or severe brain injuries than among patients with mild brain injuries (control group). The risk of developing a memory disorder was 2.5 times higher among the 41-50 age group, and 2.0 times higher among the 18-40 and 51-60 groups. The relative risk was slightly lower (1.5 times higher) among the 61-65 group, as the age-related risk for a memory disorder is already higher. Credit: R Raj and J Kaprio / University of Helsinki

More information: Raj R, Kaprio J, Korja M, Mikkonen ED, Jousilahti P, Siironen J (2017) Risk of hospitalization with neurodegenerative disease after moderate-to-severe traumatic brain

injury in the working-age population: A retrospective cohort study using the Finnish national health registries. *PLoS Med* 14(7): e1002316.
doi.org/10.1371/journal.pmed.1002316

Provided by Public Library of Science

Citation: How much does dementia risk increase after traumatic brain injury? (2017, July 5)
retrieved 6 May 2023 from
<https://medicalxpress.com/news/2017-07-traumatic-brain-injury-dementia-working-age.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.