


# Lumpectomy combined with lift a good option for women with breast cancer

October 18 2017

---



Micrograph showing a lymph node invaded by ductal breast carcinoma, with extension of the tumour beyond the lymph node. Credit: Nephron/Wikipedia

A collaborative effort between a surgical oncologist and plastic surgeon can provide an opportunity for women diagnosed with breast cancer to not only have the lump removed from the breast, but also get a breast lift during the same procedure. Dr. Sebastian Winocour, assistant professor in the Michael E. DeBakey Department of Surgery at Baylor College of Medicine who specializes in plastic surgery, says that women diagnosed with breast cancer should consider this option early, because performing this procedure after radiation has a higher rate of complications.

"When women are presented with a [breast cancer](#) diagnosis, they frequently choose between having a total mastectomy, which involves removing the breast completely, or a lumpectomy, which involves removing part of the breast and then receiving [radiation](#) afterward," said Winocour, who also is a member of the Dan L Duncan Comprehensive Cancer Center at Baylor.

If the lumpectomy option is selected, women who have breasts that sag also can inquire about having a lift during the lumpectomy procedure. During this procedure, the two surgeons work together to achieve the best cosmetic results while still fully removing the [cancer](#).

"It is an oncologically sound operation," Winocour said.

Surgeons wait for the preliminary pathology report during the surgery to confirm that all of the cancer has been removed before the plastic surgeon performs the breast lift. The procedure takes longer than a traditional lumpectomy, and once the patient heals, which is usually within four weeks, she receives radiation treatment.

The [plastic surgeon](#) and patient also can discuss whether to perform the lift on the other breast at the same time or at a later time, and there are considerations for either option. If the decision is made to lift both breasts at the same time, the surgeon will often make the other breast

smaller, because radiation will slightly shrink the affected breast. Patients also can wait until after radiation to lift the second breast, but the breasts would be uneven until the second procedure is complete.

"It's important to consider this procedure and discuss with the surgeon early on because it is difficult to have this [procedure](#) on the affected [breast](#) after radiation due to potential complications," Winocour said. "In many cases, this is a good opportunity to have this lift done."

Women often report feeling rejuvenated and youthful afterward, Winocour said, and medical insurance usually covers it.

Provided by Baylor College of Medicine

Citation: Lumpectomy combined with lift a good option for women with breast cancer (2017, October 18) retrieved 29 February 2024 from <https://medicalxpress.com/news/2017-10-lumpectomy-combined-good-option-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.