

# Sleeve gastrectomy, common weight-loss surgery, lowers women's tolerance to alcohol

November 20 2017

---

Women who have had gastric sleeve surgery to lose weight may want to consider limiting the number of alcoholic drinks they consume post-surgery.

A new study from a team of researchers at the University of Illinois and Washington University School of Medicine in St. Louis found that after undergoing sleeve gastrectomy, women could be legally intoxicated after drinking half the number of drinks than women who did not have this [surgery](#).

Sleeve gastrectomy, similar to another weight-loss surgery, Roux-en-Y gastric bypass (RYGB), speeds up alcohol absorption to the bloodstream. After drinking, [blood alcohol levels](#) increase much faster and reach higher levels than what would be expected before surgery, explains Marta Yanina Pepino, an assistant professor in the Department of Food Science and Human Nutrition at U of I who led the study.

"After having a sleeve gastrectomy, if a woman has a couple of drinks, she could be exposing her brain to [blood](#) alcohol levels that are achieved in a woman without surgery when she consumes four or five drinks," adds first author M. Belen Acevedo, a postdoc in Pepino's group at U of I. "Drinking, such that it raises blood alcohol levels above legal drinking limits, is considered a binge drinking episode and has been associated with an increased risk of developing [alcohol problems](#)."

The study is available online in the journal *Surgery for Obesity and*

### *Related Diseases.*

The researchers wanted to learn whether sleeve gastrectomy, currently the most frequently used weight-loss surgical procedure used in the United States and the world, would affect how the body handles alcohol, also known as alcohol pharmacokinetics.

"We know that changes in the pharmacokinetics of a drug are important because the faster a drug of abuse reaches your brain, the higher the potential for addiction," Pepino explains. "Although it is well known that RYGB affects how the body handles alcohol, findings from earlier studies on the effects of sleeve gastrectomy on alcohol pharmacokinetics are contradictory."

Out of three previously published studies, Pepino says that two found that sleeve surgery did not change blood alcohol levels and one study found that sleeve surgery causes a higher blood alcohol peak. "However, all previous studies on sleeve gastrectomy, estimated blood alcohol levels by using breathalyzers, which might not reliably estimate peak blood alcohol levels in bariatric patients," she adds.

For the current study, 11 women who had sleeve surgery and eight who had RYGB surgery within 1-5 years before the study began, as well as nine women who had not had either surgery, were evaluated in two sessions. In one session the women consumed the equivalent of two standard alcoholic drinks over a 10-minute period. At another, they consumed non-alcoholic drinks. At each visit, the researchers measured blood alcohol levels by using gas chromatography and compared them to estimations of blood alcohol levels obtained from breath alcohol measures using a breathalyzer. They also used a questionnaire to determine the women's level of drunkenness.

Only women were included in the study because they comprise the

majority of patients who get bariatric surgery, Pepino says.

For the women in the non-surgery group, blood alcohol contents peaked at 0.6 g/L about 26 minutes after they finished drinking. In those who had sleeve surgery and RYGB, blood alcohol contents peaked at 1.1 and 1.0 g/L about 9 and 5 minutes, respectively, after finishing the drink. The [women](#) who had sleeve surgery or RYGB also reported more intense feelings of drunkenness.

Another finding showed that breathalyzer values compared to those measured by gas chromatography, which Pepino calls the "gold-standard" technique, underestimated blood alcohol by 27 percent. Because of this finding and the fact that breathalyzers must be used 15 minutes after the final drink of alcohol, Pepino points out that breathalyzers may not be a reliable way to get accurate [alcohol](#) levels from those who have had sleeve or RYGB surgery.

"Bariatric surgeries are the most effective long-term treatment of obesity that we know of today, and findings from this study or others showing associations between RYGB and increased risk to develop alcoholism are not to say we should not perform these procedures," Pepino says. "The therapeutic effects of these surgical procedures are unparalleled. People recover from diseases such as diabetes and are able to leave many of their medications because of these procedures.

"We hope our finding motivates surgeons and the team of bariatric support professionals interacting with bariatric patients to discuss the potentially serious consequences of [moderate alcohol consumption](#) following [sleeve gastrectomy](#) and RYGB."

**More information:** María Belén Acevedo et al, Sleeve gastrectomy surgery: When two alcoholic drinks are converted to four, *Surgery for Obesity and Related Diseases* (2017). [DOI: 10.1016/j.soard.2017.11.010](https://doi.org/10.1016/j.soard.2017.11.010)

Provided by University of Illinois at Urbana-Champaign

Citation: Sleeve gastrectomy, common weight-loss surgery, lowers women's tolerance to alcohol (2017, November 20) retrieved 12 February 2024 from

<https://medicalxpress.com/news/2017-11-sleeve-gastrectomy-common-weight-loss-surgery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.