

# Acetaminophen use during pregnancy associated with elevated rate of language delay in girls

January 10 2018

---



Credit: CC0 Public Domain

In the first study of its kind, researchers from the Icahn School of Medicine at Mount Sinai found an elevated rate of language delay in girls at 30 months old born to mothers who used acetaminophen during

pregnancy, but not in boys.

This is the first study to examine [language development](#) in relation to [acetaminophen](#) levels in urine.

The study will be published online January 10 at 3:28 am EST in *European Psychiatry*.

The Swedish Environmental Longitudinal, Mother and Child, Asthma and Allergy study (SELMA) provided data for the research. Information was gathered from 754 women who were enrolled into the study in weeks 8-13 of their pregnancy. Researchers asked participants to report the number of acetaminophen tablets they had taken between conception and enrollment, and tested the acetaminophen concentration in their urine at enrollment. The frequency of [language delay](#), defined as the use of fewer than 50 words, was measured by both a nurse's assessment and a follow-up questionnaire filled out by participants about their child's [language](#) milestones at 30 months.

Acetaminophen was used by 59 percent of the women in early pregnancy. Acetaminophen use was quantified in two ways: High use vs. no use analysis used women who did not report any use as the comparison group. For the [urine analysis](#), the top quartile of exposure was compared to the lowest quartile.

Language delay was seen in 10 percent of all the children in the study, with greater delays in boys than girls overall. However, girls born to mothers with higher exposure—those who took acetaminophen more than six times in early pregnancy—were nearly six times more likely to have language delay than girls born to mothers who did not take acetaminophen. These results are consistent with studies reporting decreased IQ and increased communication problems in children born to mothers who used more acetaminophen during pregnancy.

Both the number of tablets and concentration in urine were associated with a significant increase in language [delay](#) in girls, and a slight but not significant decrease in boys. Overall, the results suggest that acetaminophen use in pregnancy results in a loss of the well-recognized female advantage in language development in early childhood.

The SELMA study will follow the children and re-examine language development at seven years.

Acetaminophen, also known as paracetamol, is the active ingredient in Tylenol and hundreds of over-the-counter and prescription medicines. It is commonly prescribed during pregnancy to relieve pain and fever. An estimated 65 percent of [pregnant women](#) in the United States use the drug, according to the U.S. Centers for Disease Control and Prevention.?

"Given the prevalence of prenatal acetaminophen use and the importance of language development, our findings, if replicated, suggest that pregnant women should limit their use of this analgesic during [pregnancy](#)," said the study's senior author, Shanna Swan, PhD, Professor of Environmental and Public Health at the Icahn School of Medicine at Mount Sinai. "It's important for us to look at language development because it has shown to be predictive of other neurodevelopmental problems in children."

Provided by The Mount Sinai Hospital

Citation: Acetaminophen use during pregnancy associated with elevated rate of language delay in girls (2018, January 10) retrieved 11 July 2023 from <https://medicalxpress.com/news/2018-01-acetaminophen-pregnancy-elevated-language-girls.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.