


Physicists link specific iron forms to Alzheimer's

May 8 2018



Credit: Leiden Institute of Physics


There have been indications for decades that there is a link between increased iron levels in the brain and Alzheimer's disease. Leiden physicists now report a distinction between different forms of iron, identifying specific iron forms that increase in Alzheimer's patients. The research is published in *Scientific Reports*.

Iron plays an important role in biological processes, including those in the brain. With cerebral disorders such as Alzheimer's disease, there are indications that the iron balance is disturbed, because researchers measure increased [iron content](#). To measure iron, scientists use an MRI

scanner, which detects iron content indirectly.

Lucia Bossoni from the Leiden Institute of Physics and the Leiden University Medical Center and her colleagues have now combined MRI with EPR and SQUID magnetometry. This enables them to distinguish different shapes in which iron occurs. They confirm previous studies that establish a link between increased iron content and Alzheimer's disease, while making this connection specific for individual iron forms. For example, the iron concentration in the mineral ferrihydrite is higher in the group of Alzheimer's patients than in the control group (see figure).

In addition, Bossoni spots a trend where certain iron forms correlate with the Braak stage. The iron concentration in the protein ferritin and the magnetic moment of the mineral magnetite appear to increase as Alzheimer's disease reaches a more advanced stage. Bossoni says, "When we talk about iron increase in connection with neurodegenerative diseases, we have to pay extra attention to the specific [iron](#) form that really matters."



Difference between the group of Alzheimer's patients (red) and the control group (blue) in the magnetic moment of magnetite (left) and the iron concentration in ferrihydrite (right). Both are generally larger in the group of Alzheimer's patients. Credit: Leiden Institute of Physics

More information: Marjolein Bulk et al. Quantitative comparison of different iron forms in the temporal cortex of Alzheimer patients and control subjects, *Scientific Reports* (2018). [DOI: 10.1038/s41598-018-25021-7](https://doi.org/10.1038/s41598-018-25021-7)

Provided by Leiden Institute of Physics

Citation: Physicists link specific iron forms to Alzheimer's (2018, May 8) retrieved 3 February 2024 from <https://medicalxpress.com/news/2018-05-physicists-link-specific-iron-alzheimer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.