

Increased adiposity tied to lower premenopausal breast CA risk

June 25 2018



(HealthDay)—Across body mass index (BMI) distribution, increased

BMI is associated with reduced risk of premenopausal breast cancer, according to a study published online June 21 in *JAMA Oncology*.

Minouk J. Schoemaker, Ph.D., from The Institute of Cancer Research in London, and colleagues from the Premenopausal Breast Cancer Collaborative Group examined the correlation of BMI with premenopausal breast cancer risk in a multicenter analysis using pooled individual-level data from 758,592 premenopausal women from 19 prospective cohorts. Participants were followed for a median of 9.3 years per participant to estimate the hazard ratios of premenopausal breast cancer in association with BMI; there were 13,082 incident cases of breast cancer during follow-up.

The researchers found that inverse linear associations of BMI with breast cancer risk were identified that were stronger for BMI at ages 18 to 24 years than for BMI at ages 45 to 54 years (hazard ratios per 5 kg/m² difference, 0.77 and 0.88, respectively). Even among non-overweight women, these inverse associations were observed. At ages 18 to 24 years there was a 4.2-fold risk gradient between the highest and lowest BMI categories (BMI \geq 35.0 vs

Citation: Increased adiposity tied to lower premenopausal breast CA risk (2018, June 25) retrieved 19 November 2023 from <https://medicalxpress.com/news/2018-06-adiposity-tied-premenopausal-breast-ca.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.