

Cancer treatments may affect cognitive function by accelerating biological aging

November 26 2018



Credit: CC0 Public Domain

Cancer treatments are suspected to accelerate certain aging processes in the body. A new study has found that indicators of such biological aging correlate with declines in cognitive function in women who had

undergone breast cancer treatment several years earlier. Published early online in *Cancer*, a peer-reviewed journal of the American Cancer Society, the findings point to an aging-like effect of cancer treatments and further connect this to cognitive decline.

Treatments for breast [cancer](#) increase patients' risks for long-term and late toxicities, including persistent fatigue, pain, and cognitive dysfunction. Certain treatments, including radiation and some chemotherapeutic drugs, work by damaging the DNA of cancer cells, but they can also cause damage to DNA of normal cells, which can contribute to accelerated biological aging.

To examine whether indicators of biological aging are related to cognitive function in breast cancer survivors, Judith E. Carroll, Ph.D., Associate Professor of Psychiatry, of the UCLA Cousins Center for Psychoneuroimmunology and the Semel Institute for Neuroscience and Human Behavior, and her colleagues evaluated a group of 94 women who had been treated for breast cancer three to six years earlier. The indicators of biological aging included elevated levels of DNA damage, reduced [telomerase enzymatic activity](#), and shorter telomere length in certain blood cells. (Telomerase is an enzyme that is important for maintaining the length of telomeres, repeat sequences of DNA at the ends of chromosomes that help maintain the health of cells and serve as a marker of cell age.)

The team found that women who had previously been treated for breast cancer who had both higher DNA damage and lower telomerase activity had lower executive function scores. In addition, lower telomerase activity was associated with worse attention and motor speed. Telomere length was not related to any of the neurocognitive domains.

"These findings are important because they provide further information about what might be happening after [cancer treatment](#) that impacts

cognitive decline in some individuals. This information can inform future research and may lead to new interventions to prevent these cognitive declines," said Dr. Carroll, who is also a member of the UCLA Jonsson Comprehensive Cancer Center. "The work is novel by identifying key factors in biological aging and connecting them to cognitive function, which initiates new avenues of research."

More information: Judith E. Carroll et al. Cognitive performance in survivors of breast cancer and markers of biological aging, *Cancer* (2018). DOI: [10.1002/cncr.31777](https://doi.org/10.1002/cncr.31777)

Provided by Wiley

Citation: Cancer treatments may affect cognitive function by accelerating biological aging (2018, November 26) retrieved 6 April 2024 from <https://medicalxpress.com/news/2018-11-cancer-treatments-affect-cognitive-function.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.