

Researcher finds gender disparities in admissions, treatment for heart attack patients, earns nomination for AHA award

November 19 2018

A new study published in the American Heart Association (AHA) journal, *Circulation*, shows that that the dramatic decrease in mortality from heart attack in recent decades is not evident in younger age groups, especially younger women. The study, led by UNC School of Medicine cardiology fellow Sameer Arora, MD, looked at data collected by the Atherosclerosis Risk in Community (ARIC) study, a hospital surveillance of acute myocardial infarction (AMI) in patients aged 35-74 years old in four U.S. cities. Analysis for Arora's study was limited to nearly 9,000 patients aged 35-54 years old from 1995 to 2014.

The overall incident of AMI, or heart attack, has decreased in recent decades in the U.S. However, when looking specifically at the 35-54 year-old age group, Arora and his colleagues discovered that this age group had an increase in heart attacks, and <u>women</u> had the largest increase of annual hospital admission due to heart attack. The proportion of heart attacks attributable to young women increased from 21 to 31 percent over 19 years.

"Heart attacks are traditionally considered an older man's disease," said Arora. "But we discovered that the proportion overall is shifting towards <u>younger women</u>. Young women are catching up to young men in <u>heart</u> attack incidence."

Arora also found that <u>young women</u> are not experiencing the same



treatment as young men. Women were less likely to receive guidelinebased AMI treatment, such as lipid-lowering therapies, non-aspirin antiplatelets, beta blockers, coronary angiography, and coronary revascularization. Women were also more likely than men in this age group to experience comorbidities, such as hypertension, diabetes, and chronic kidney disease.

"Young women are not being treated the same way as young men," Arora said. "This indicates a perception problem; physicians possibly are not believing that <u>heart attack</u> is as big of an issue for women. There should be a different set of treatment guidelines for women, and their unique risk factors should be recognized as well."

The study's senior author is Melissa Caughey, Ph.D., research instructor of medicine in the division of cardiology, who also served as Arora's faculty mentor. UNC's Rick Stouffer, MD, chief of cardiology, along with Wayne Rosamond, Ph.D., and Anna Kucharska-Newton, MPH, Ph.D., both professors in the department of epidemiology in the Gillings Schools of Global Public Health, were also authors.

More information: Twenty Year Trends and Sex Differences in Young Adults Hospitalized with Acute Myocardial Infarction: The ARIC Community Surveillance Study, *Circulation*, <u>DOI:</u> <u>10.1161/CIRCULATIONAHA.118.037137</u>

Provided by University of North Carolina at Chapel Hill School of Medicine

Citation: Researcher finds gender disparities in admissions, treatment for heart attack patients, earns nomination for AHA award (2018, November 19) retrieved 20 November 2023 from <u>https://medicalxpress.com/news/2018-11-gender-disparities-admissions-treatment-heart.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.