

Strong painkillers increase the risk of hip fracture among persons with Alzheimer's disease

November 26 2018



Credit: CC0 Public Domain

People using strong painkillers, opioids, have twice the risk of hip fracture compared to non-opioid users, a new study from the University

of Eastern Finland shows. The risk was highest in the first two months of opioid use. The results were published in the *PAIN* journal.

The risk of hip fracture increased with increasing opioid strength. Use of weak opioids, such as codeine and tramadol, was not associated with the risk of hip fracture. However, moderately strong opioids, such as buprenorphine, were associated with a two-fold risk, and strong opioids, such as oxycodone and [fentanyl](#), were associated with almost a three-fold risk, compared with non-use. Buprenorphine and fentanyl were almost solely used as patches in this study.

Opioids are used for the treatment of severe pain in cases where other [painkillers](#) fail to produce a sufficient response. Opioids impact on the central nervous system and for this reason, opioid use may cause attentional impairment or drowsiness. Opioids are known to increase the risk of falls which, in turn, may lead to hip fracture in [older people](#).

The study was based on the nationwide register-based MEDALZ study conducted at the University of Eastern Finland. For this study, 23,100 community-dwelling persons diagnosed with Alzheimer's disease in Finland during 2010-2011 were included. From this sample, new opioid users were identified and matched with [opioid](#) non-users.

More information: Heidi Taipale et al. Incident opioid use and risk of hip fracture among persons with Alzheimer disease, *PAIN* (2018). [DOI: 10.1097/j.pain.0000000000001412](https://doi.org/10.1097/j.pain.0000000000001412)

Provided by University of Eastern Finland

Citation: Strong painkillers increase the risk of hip fracture among persons with Alzheimer's disease (2018, November 26) retrieved 19 November 2023 from

<https://medicalxpress.com/news/2018-11-strong-painkillers-hip-fracture-persons.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.