

Risk factors ID'd for site infection after orthopedic surgery in seniors

February 28 2019



(HealthDay)—Five risk factors are independently associated with

surgical site infection among geriatric patients undergoing elective orthopedic surgery, according to a study published online Feb. 20 in the *International Wound Journal*.

Zhiquan Liang, from the First Hospital of Jilin University in Changchun, China, and colleagues retrospectively reviewed [medical records](#) (January 2014 through September 2017) for 4,818 patients ≥ 60 years old undergoing elective orthopedic surgeries in order to assess the risk for SSI.

The researchers found that within one postoperative year, 74 patients developed SSIs. The overall incidence of SSI was 3.64 percent, with 0.4 percent for deep infection and 1.1 percent for superficial infection. The most common causative pathogens were *Staphylococcus aureus* (53.2 percent) and coagulase-negative *staphylococci* (23.4 percent). Nearly half of *S. aureus* SSIs (12 of 25) were caused by methicillin-resistant *Staphylococcus aureus*. Diabetes mellitus (odds ratio [OR], 3.7), morbid obesity (OR, 2.6), tobacco smoking (OR, 4.2), surgical duration >75 th percentile (OR, 1.9), and albumin

Citation: Risk factors ID'd for site infection after orthopedic surgery in seniors (2019, February 28) retrieved 20 November 2023 from <https://medicalxpress.com/news/2019-02-factors-idd-site-infection-orthopedic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.