


Diattenuation imaging—a promising imaging technique for brain research

March 20 2019




Diattenuation Imaging (DI) provides structural information about brain tissue. The colors reveals the direction of polarization for which a maximum amount of light passes through the brain section. Regions for which this polarization direction runs parallel (or perpendicular) to the fiber direction are marked in green (or magenta). Credit: Miriam Menzel et al., *Scientific Reports* (2019), DOI:10.1038/s41598-019-38506-w (CC BY 4.0)

A new imaging method provides structural information about brain tissue that was previously difficult to access. Diattenuation Imaging (DI), developed by scientists at Forschungszentrum Jülich and the University of Groningen, allows researchers to differentiate regions with many thin nerve fibres from regions with few thick nerve fibres. With current imaging methods, these tissue types cannot easily be distinguished.

The DI method is based on 3-D polarized light imaging (3-D-PLI), a [neuroimaging technique](#) developed at Forschungszentrum Jülich, which reveals [nerve](#) fibre pathways with micrometre resolution. The technique is used, for instance, in the European Human Brain Project to investigate the 3-D [fibre](#) structures of the [brain](#) in unprecedented detail.

During a 3-D-PLI measurement, histological brain sections are illuminated with polarized light. Depending on how the direction of oscillation (polarization) is oriented relative to the nerve fibres, the light is refracted to different degrees, allowing the computation of the spatial orientation of the nerve fibres. This effect, called birefringence, is mainly caused by the myelin sheath, an insulating layer that surrounds many nerve fibres in the brain.

While 3-D-PLI measures the polarization-dependent refraction of light, a diattenuation measurement determines the polarization-dependent attenuation of light, i.e. how much the intensity of polarized light is reduced when passing through the brain section. The measurement is performed with the same apparatus as 3-D-PLI, whereby two filters are removed.




How the new imaging method Diattenuation Imaging works: 3D-PLI (left) reveals the direction of the nerve fibers. A diattenuation measurement (right) reveals the direction of polarization for which a maximum amount of light passes through the brain section. Regions for which this polarization direction runs parallel (or perpendicular) to the fibre direction are marked in green (or magenta). Credit: Forschungszentrum Jülich / Miriam Menzel, Tobias Schlößer

The scientists discovered that diattenuation imaging—a combined measurement of diattenuation and 3-D-PLI—can distinguish between different brain regions. In some regions, the [brain tissue](#) is maximally transparent when the polarization of the light is oriented parallel to the nerve fibres. In other regions, the [tissue](#) is maximally transparent when

the polarization is oriented perpendicularly to the nerve fibres. How the tissue behaves depends, among other things, on the time after embedding the brain sections.

Using simulations on the former Jülich supercomputer JUQUEEN, the researchers could show that the observed effects also depend on other tissue properties like the diameter of the fibres or the thickness of the myelin sheaths. This makes diattenuation imaging a valuable extension to 3-D-PLI, enabling a more precise investigation of brain tissue. In the future, the DI method could be used to study [neurodegenerative diseases](#) like multiple sclerosis or multisystem atrophy (MSA), which go along with alterations of the myelin sheath. In addition, the technology helps to make pathological changes visible and to identify connected regions and tissue types, assisting the complex reconstruction of the brain.



The regional differences that become apparent with Diattenuation Imaging depend, among other things, on the time after embedding the brain sections (left: freshly measured brain section, right: seven weeks later). Credit: Miriam Menzel et al., *Scientific Reports* (2019), DOI:10.1038/s41598-019-38506-w (CC BY 4.0)

More information: Miriam Menzel et al, Diattenuation Imaging reveals different brain tissue properties, *Scientific Reports* (2019). [DOI: 10.1038/s41598-019-38506-w](https://doi.org/10.1038/s41598-019-38506-w)

Provided by Forschungszentrum Juelich

Citation: Diattenuation imaging—a promising imaging technique for brain research (2019, March 20) retrieved 5 February 2024 from <https://medicalxpress.com/news/2019-03-diattenuation-imaginga-imaging-technique-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.