


If you're coming off antidepressants, withdrawals and setbacks may be part of the process

April 12 2019, by Michael Musker



Credit: AI-generated image

Depression is a common mental disorder affecting over [300 million people across the world](#). It's estimated that [one in ten](#) people in Australia (10.4%) suffer from depression.

Alongside other therapies, medication is often used to treat patients with depression. About [8% of Australians](#) are taking [antidepressants](#) for depression, anxiety and related conditions (but in this article we're focusing on depression).

Antidepressants are powerful drugs that affect the way the neurotransmitters in our brains work, usually with the outcome of increasing the overall amount of a chemical called [serotonin](#). Low levels of serotonin cause low mood, so increasing the availability of this transmitter improves mood.

To be in a position where you feel ready to come off antidepressants is a good thing. But it's not necessarily an easy thing – particularly if you've been taking antidepressants for a long time.

There is a possibility of withdrawal symptoms, and even the chance of relapsing back into a depression. These risks can't be removed, but they can be managed.

The illness trajectory

Depression can last for a minimum of two weeks, but will more commonly last for months or years. For some, it will be present for the duration of their lives.

Typically, those who suffer from a major depressive disorder will be taking antidepressants for many years.

Antidepressants can sometimes cause [side effects](#) such as weight gain, fatigue, sexual dysfunction and insomnia. It's understandable that people will want to stop taking these drugs as soon as possible, or after they have felt well for a considerable time.

But this process carries risks, so it's important to establish a plan with your doctor before proceeding.

Antidepressant discontinuation syndrome

Your body has become used to having the drugs in its system, so withdrawal can cause side effects.

A condition called [antidepressant discontinuation syndrome](#) affects about 20% of people withdrawing from antidepressants.

This syndrome can cause flu-like symptoms (headache, body aches and sweating), difficulty sleeping, irritability, feeling sick (nausea or even vomiting), disturbance in balance, confusion, anxiety and agitation.

These symptoms may only last for a week, but could have more serious consequences if not monitored properly, such as falling back into depression. It's important to have a withdrawal plan with your GP or prescribing doctor, so that these symptoms can be measured and medications adjusted accordingly.



Credit: Константин from Pexels

Some antidepressants such as Paroxetine and Venlafaxine ([selective serotonin reuptake inhibitors](#)) are [more likely to cause withdrawal](#), so extra care should be taken with these medicines. Other drugs like Fluoxetine can be withdrawn over a shorter period, or even act as a buffer while withdrawing from other antidepressants.

Steps to take if you're coming off antidepressants

Coming off antidepressants too quickly may increase the risk that a

person will fall into a depression again and need to revert to taking medication, creating a cycle of depression.

The [American Psychiatric Association](#) and the [UK National Collaborating Centre for Mental Health](#) suggest antidepressants are taken for up to several weeks after the depression symptoms have ceased, particularly if you have been on them for many years. The key aim here is to prevent a relapse into depression.

It takes between four to eight weeks for an antidepressant to start working effectively, and it will take just as long for your body to get used to not being on them.

A general guide is to withdraw from antidepressants over around [four weeks](#), but longer if you are on a group of drugs called monoamine oxidase inhibitors.

This process involves a gradual reduction of the amount of the drug you're taking over time, according to the plan you and your doctor have agreed upon. Keeping track of how this is affecting your mood is helpful.

A [systematic review](#) found several interventions helpful when thinking about stopping antidepressants. The most effective was a form of psychological treatment called [cognitive behavioural therapy](#) (CBT).

CBT is a talking therapy that is generally provided in hour-long sessions over a period of about 16 weeks during and after the withdrawal phase.

CBT can be combined with an activity called [mindfulness](#), a type of relaxation that helps you focus on your thoughts.

What if you do experience withdrawal?

If you find yourself suffering from the symptoms of withdrawal, returning to a lower dose of medication and slowing down the discontinuation over a longer period may be a sensible way to go.

You can also focus on some [key activities](#) known to have a positive effect upon [depression](#). These might include forming good sleep habits, increasing the amount of exercise you do, engaging in your favourite hobbies or social activities, or adopting a healthy low carbohydrate diet.

The [Black Dog Institute](#) recommend you monitor your mood in a [mood chart](#), which will give you a clear picture of how you're tracking. This enables you to see the danger points when your mood falls below par, and consider when you may need to seek help.

There are many [free interactive resources](#) online which may be helpful, alongside professional medical care, as you navigate coming off [antidepressants](#). You can also focus on [positive mental health strategies](#) that build your well-being and resilience.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: If you're coming off antidepressants, withdrawals and setbacks may be part of the process (2019, April 12) retrieved 21 November 2023 from <https://medicalxpress.com/news/2019-04-youre-antidepressants-setbacks.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.