


Cell-cell signals in developing heart

June 14 2019, by Leigh Macmillan



Human heart. Credit: copyright American Heart Association

During late stages of heart development, interactions between the endocardium (the inner layer of cells) and the myocardium (the heart muscle) are known to be crucial. Signaling between these two cell layers during the earliest stages of heart development has been more difficult to study.

Now, H. Scott Baldwin, MD, and colleagues have generated a model to explore endocardial-myocardial interactions in vitro. They found that eliminating endocardial cells reduced the number of contracting myocardial cells and the expression of myocardial-specific genes. Addition of the growth factor Bmp2 partially rescued myocardial cell maturation and function.

The studies, reported in the journal *Development*, provide direct evidence of a role for the endocardium during early [heart muscle](#) development and could guide stem cell-based therapies for generating myocardium.

The findings also point to the possibility that altered endocardial-myocardial interactions during early development may contribute to congenital heart defects, which occur in nearly 1% of [live births](#) and remain the leading cause of death due to birth defects.

More information: Leshana Saint-Jean et al. Myocardial differentiation is dependent upon endocardial signaling during early cardiogenesis in vitro, *Development* (2019). [DOI: 10.1242/dev.172619](https://doi.org/10.1242/dev.172619)

Provided by Vanderbilt University

Citation: Cell-cell signals in developing heart (2019, June 14) retrieved 19 April 2023 from <https://medicalxpress.com/news/2019-06-cell-cell-heart.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--