


# Model projects sharp rise in alcohol-related liver disease

June 4 2020, by Brian Burns

---



Credit: CC0 Public Domain

Alcohol-related liver disease is currently the most common reason for liver transplantation in the United States. In recent years, high-risk drinking of alcohol—defined as exceeding daily drinking limits (more than four or five standard drinks for women and men, respectively) at least weekly for a whole year—has increased in nearly all sociodemographic groups, especially in women.

A new analysis by researchers at Massachusetts General Hospital, Harvard Medical School and Georgia Tech indicates that cases of [alcohol-related liver disease](#) will rise dramatically in the coming years without drastic steps to reduce high-risk drinking rates.

To project the impact of alcohol-related [liver](#) disease over the next two decades, the team developed a model of drinking patterns and alcohol-related liver disease in individuals born in 1900 to 2012 in the United States, with projections up to 2040. The model relied on data from multiple sources: the National Epidemiologic Survey on Alcohol and Related Conditions, the National Institute of Alcohol Abuse and Alcoholism, the US National Death Index, the National Vital Statistics System and numerous published studies. The model was validated in that it closely reproduced the trends in deaths due to alcohol-related liver disease that were observed from 2005 to 2018.

In *The Lancet Public Health* analysis, future trends in alcohol-related liver disease were modelled under three potential scenarios based on the level of interventions to address high-risk drinking:

- Without any changes in trends in alcohol consumption (status quo), more than 1 million people could die from alcohol-related

liver disease by 2040, of whom 35% are projected to be younger than 55 years

- Reducing high-risk drinking rates to 2001 levels could prevent 35,000 deaths during the same time period
- In contrast, decreasing the high-risk drinking rate by 3.5% per year—similar to the decrease in the rate of tobacco consumption observed after implementation of policies and social interventions in the 1960s to reduce smoking in the U.S.—could prevent 299,000 deaths compared with the status quo scenario (a 30% decrease)

The analysis also projected that in comparison with status quo, decreasing the high-risk drinking rates by 3.5% per year could prevent alcohol-related liver cancer and decompensated cirrhosis (a form of advanced liver disease) by 30% from 2019 to 2040.

"Our study underscores the need to bring alcohol-related disease to the forefront of policy discussions and identify effective policies to reduce high-risk drinking in the U.S.," said senior author Jagpreet Chhatwal, Ph.D., a senior scientist at the MGH Institute for Technology Assessment and an assistant professor at Harvard Medical School.

Lead author Jovan Julien, MS, a Ph.D. student at Georgia Tech, noted that the impact of high-risk drinking takes years to observe. "Our model highlights the long-term risk, especially for younger generations whose drinking continues to outpace older generations," he said. "Given the practical infeasibility of clinical trials to study long-term effects of increased [alcohol consumption](#), data-driven modeling studies such as ours are valuable for understanding the implications of high-risk [drinking](#) and making informed policy decisions," added co-author Turgay Ayer, Ph.D., MSc, an associate professor at Georgia Tech.

**More information:** Jovan Julien et al. Projected prevalence and

mortality associated with alcohol-related liver disease in the USA, 2019–40: a modelling study, *The Lancet Public Health* (2020). [DOI: 10.1016/S2468-2667\(20\)30062-1](https://doi.org/10.1016/S2468-2667(20)30062-1)

Provided by Massachusetts General Hospital

Citation: Model projects sharp rise in alcohol-related liver disease (2020, June 4) retrieved 11 June 2024 from

<https://medicalxpress.com/news/2020-06-sharp-alcohol-related-liver-disease.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.