

Women taking menopausal hormone therapy may be more resistant to urine infections

July 20 2020



Credit: CC0 Public Domain

In the first analysis of its kind, U.S.-based doctors have shown that women who take menopausal hormone therapy (MHT, also known as HRT) have a greater variety of beneficial bacteria in their urine, possibly

creating conditions that discourage urinary tract infections. The study also shows that women who suffer from recurrent infections have a lower diversity of bacteria in their urine, making them more prone to infections. This work is presented at the European Association of Urology Virtual Congress.

Around 50 to 60% of women will suffer from a painful urinary tract infection (UTI) during their lifetime, with women being around 8x more likely than men to suffer a UTI. Urine is not sterile, and normal [urine](#) contains a variety of bacteria, fungi and viruses. Increasingly, doctors are finding that the people's bodies act as host to a variety of harmless resident bacteria, which may help in processes such as digestion.

Now the first controlled, systematic analysis of the diversity of bacteria in the urine of postmenopausal women has shed light on the relationship between bacterial diversity and recurrent urinary infections and susceptibility.

A group led by Dr. Nicole J. De Nisco, assistant professor, University of Texas at Dallas, collected and analyzed bacterial DNA from the urine of 75 postmenopausal women who were attending the Urology Clinic at UT Southwestern Medical Center in Dallas, Texas. They were able to divide the women into three groups: 25 women who had never had a UTI, 25 women who had recurrent infections in the past (i.e., three or more infections per year) but no current infection, and 25 women who had recurrent infections in the past and were currently infected. Recurrent UTI is a growing clinical concern that disproportionately affects postmenopausal women.

Dr. De Nisco says, "The DNA analysis showed two things. Firstly, women who have recurrent infections have fewer types of bacteria in their urine than women who do not have infections; women who don't have recurrent UTIs have around 10 times greater variety of bacteria.

Secondly, 34 of the women were taking [menopausal hormone therapy](#), and they tended to have more Lactobacillus-type bacteria in their urine, which may imply that the estrogen in MHT supports the growth of Lactobacillus in the urogenital tract. We also found that women who were taking MHT via patches or orally had more Lactobacillus than women taking MHT via vaginal cream. This is important, because Lactobacilli are known to play a protective role against infections in the vagina and this function may be conserved in the urogenital tract. If Lactobacillus really can help with UTIs, development of a probiotic to use in conjunction with MHT may avoid [antibiotic use](#) in these women, but this needs to be tested. This finding also fits with the observation that postmenopausal women, who have less natural estrogen, tend to have more infections."

She continued, "As far as we know, this is the first focused analysis of the [bacteria](#) in the urogenital tract of postmenopausal women, and the results indicate that estrogen use is associated with high levels of Lactobacilli in the urine, which could possibly offer some protection against infection. However, this is a small study and needs to be repeated with a larger group of women. We now need to look more closely at whether managing the bacterial content of the urogenital system might affect susceptibility to infection. Additional studies need to be performed to determine a causal link between estrogen therapy, Lactobacillus colonization and prevention of UTI. Possible future applications may include administering probiotic vaginal pessaries to women at risk of infection and perhaps prescribing estrogen therapy. However it's too early to bring these into clinical practice. We know that estrogen therapy is not recommended for all women going through menopause, and the optimal formulations of probiotic pessaries need to be developed and their use in [women](#) with UTI would need to be tested. This is the next step in our research."

Professor Florian Wagenlehner (Giessen, Germany), chair of the EAU

specialist group on Infections in Urology, who was not involved in the study, said, "This study, which evaluates urine microbiome in [postmenopausal women](#), shows that patients with fewer urinary infections tend to have a greater species diversity in their urine. We see more and more evidence that diversity in the microbiome prevents infections. The growth of lactobacillus species is dependent on estrogen, so estrogen supplementation may be a helpful therapy in postmenopausal patients."

Provided by European Association of Urology

Citation: Women taking menopausal hormone therapy may be more resistant to urine infections (2020, July 20) retrieved 14 February 2024 from <https://medicalxpress.com/news/2020-07-women-menopausal-hormone-therapy-resistant.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.