

# Antibiotic overuse reduces urothelial cancer survival rates

August 17 2020

---



Credit: CC0 Public Domain

The over-prescribing of antibiotics is reducing survival rates in patients with urothelial carcinoma and it needs to be stopped to avoid the serious risk posed by resistant bacterial infections, according to a new medical study.

By analyzing data collected within [clinical trials](#) on a common immunotherapy treatment called atezolizumab, cancer researchers at Flinders University have found [antibiotics](#) are consistently associated with worse [survival rates](#) in patients with [urothelial carcinoma](#).

Clinical Pharmacology Dr. Ashley Hopkins, from the Precision Medicines Group, says the study's findings suggest antibiotics may specifically reduce the effectiveness of cancer immunotherapies.

"We demonstrated that antibiotic use is directly associated with worse survival in patients with urothelial carcinoma when they're treated with atezolizumab. But no antibiotic association was observed in patients treated with chemotherapy, suggesting that antibiotics may specifically reduce the effectiveness of cancer immunotherapies," says Dr. Hopkins.

Previous research suggests up to 50% of [antibiotic use](#) in cancer treatments is prescribed inappropriately as a result of false misconceptions about there being no consequences, but this new study in the *European Urology* journal demonstrates that over reliance needs to stop.

"There are concerns that practitioners are over prescribing antibiotics like atezolizumab, potentially increasing the risk of resistant bacterial infections, and this study's findings suggest a need for extra caution when an immune checkpoint inhibitor (ICI) like this is being used."

The authors say future research will need to explore the effect of antibiotics on other immune checkpoint inhibitors (ICI) and confirm whether ICI's remain the treatment of choice in cancer patients requiring antibiotics.

"These results provide strong justification for prospective studies to tease out whether antibiotics are primarily a surrogate of an unfit or

immunodeficient patient or whether antibiotic effects on the gut microbiota are having casual impacts on ICI efficacy," says Dr. Hopkins.

"If the latter is true, in patients at a high risk of recurrent infections, it may need to be considered whether ICI therapy is the most appropriate way to go."

**More information:** Ashley M. Hopkins et al. Concomitant Antibiotic Use and Survival in Urothelial Carcinoma Treated with Atezolizumab, *European Urology* (2020). [DOI: 10.1016/j.eururo.2020.06.061](https://doi.org/10.1016/j.eururo.2020.06.061)

Provided by Flinders University

Citation: Antibiotic overuse reduces urothelial cancer survival rates (2020, August 17) retrieved 3 February 2024 from <https://medicalxpress.com/news/2020-08-antibiotic-overuse-urothelial-cancer-survival.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.