

Researchers identify a potentially safer approach to opioid drug development

August 24 2021, by Kimberly P. Cummings



Credit: Unsplash/CC0 Public Domain

Opioids are powerful painkillers, but their use is hindered because patients become tolerant to them, requiring higher and higher doses, and overdoses can cause respiratory depression and death. A recent study

from researchers at the UC Davis Center for Neuroscience contradicts existing thinking about how opioid drugs cause tolerance and respiratory depression, and suggests a new, balanced approach to developing safer analgesics. The work was published July 13 in *Neuropsychopharmacology*.

"The holy grail of [opioid](#) research is to determine the ideal properties of an opioid analgesic for maximizing pain relief while reducing the [adverse side effects](#)," said Jennifer Whistler, senior author on the paper and professor of physiology and membrane biology in the UC Davis School of Medicine. "This goal has become even more urgent in light of the devastation wreaked by the opioid overdose crises and the failure to identify other non-opioid targets for the treatment of severe and persistent pain."

Whistler, who is associate director of the UC Davis Center for Neuroscience, has been researching the neurobiology of addictive disorders and their comorbidities and how to make safer opioids for more than 20 years.

Searching for new opioids with fewer side effects

Opioid drugs work by connecting to the mu opioid receptor (MOR) on cells. This receptor in turn signals through G-protein and can also engage a protein called arrestin-3. The prevailing view has been that engagement of the mu opioid receptor with arrestin-3 is responsible for the two treatment-limiting side effects of opioids: The respiratory depressive effects that cause overdose death and the development of analgesic tolerance that leads to dose escalation and increased risk of addiction and overdose death.

This doctrine has led to an almost two decades-long, highly visible search for new "ultra G protein biased" opioids that potently activate G

protein but do not engage arrestins.

It has also led to the investment of millions of dollars into the clinical development of these new "ultra-biased" opioids, including recently FDA-approved Oliceridine, which Whistler predicts will have a higher liability to produce tolerance and addiction than our existing opioid therapeutics.

"Contrary to the prevailing hypothesis, we have found that arrestin-3 engagement prevents analgesic tolerance and does not exacerbate [respiratory depression](#)," said Whistler. "We used a powerful combination of genetic and pharmacological approaches to demonstrate this point."

The Whistler Lab team challenged the prevailing hypothesis with a panel of six clinically relevant [opioid drugs](#) and mice of three distinct genotypes with varying abilities to promote morphine-mediated arrestin-3 engagement. With this genetic and pharmacological approach, they showed that arrestin-3 recruitment does not promote respiratory depression and that effective arrestin-3 engagement reduced, rather than exacerbated, the development of analgesic tolerance.

New approach to developing opioid drugs

Whistler's data suggest an entirely new approach to the development of opioid therapeutics.

"Specifically, we propose a shift in effort to develop "balanced" opioid analgesics that strongly promote arrestin-3 engagement, much like our endogenous endorphins do," said Whistler. "In light of both the pressing need for new analgesics and the paradigm-shifting nature of our findings, we believe the time has come to try this new approach."

These studies suggest that future development of safer opioids should

focus on identifying such "balanced" opioid ligands that recruit both G protein and arrestin-3, thereby mimicking the signaling profile of most endogenous mu-opioid receptor agonists.

"There are a plethora of biased agonists, including all the opioids we take for pain. We cannot know whether a balanced approach will lead to safer opioids, until we have a library of such molecules to test," Whistler said.

More information: Li He et al, Pharmacological and genetic manipulations at the μ -opioid receptor reveal arrestin-3 engagement limits analgesic tolerance and does not exacerbate respiratory depression in mice, *Neuropsychopharmacology* (2021). [DOI: 10.1038/s41386-021-01054-x](https://doi.org/10.1038/s41386-021-01054-x)

Provided by UC Davis

Citation: Researchers identify a potentially safer approach to opioid drug development (2021, August 24) retrieved 15 March 2023 from <https://medicalxpress.com/news/2021-08-potentially-safer-approach-opioid-drug.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
