

Cancer survival rates are improving in Australia

November 30 2021



Credit: Unsplash/CC0 Public Domain

Cancer survival rates in Australia are continuing to improve and the rate at which Australians are being diagnosed with cancer has been declining since 2008, according to a new report by the Australian Institute of



Health and Welfare (AIHW).

The report, "Cancer in Australia 2021," shows a downward trend in the rate of new cancer diagnoses since 2008, but an increasing rate for females, albeit from a lower base than males.

"The estimates for 2021 presented in the report are based on actual data up to the end of 2017. They don't take into account potential health service disruptions due to COVID-19, but are still useful for identifying trends in cancer cases in Australia," AIHW spokesperson Justin Harvey said.

"It is estimated that around 151,000 new cases of cancer will be diagnosed in Australia by the end of 2021, which is an increase from around 47,500 cases in 1982. However, there has been a 5% decrease in the incidence rate over recent years from the peak of 508 cases per 100,000 people in 2008 to 486 cases per 100,000 people in 2021."

The report shows that five-year survival rates from all cancers combined had improved from 51% in 1988–1992 to 70% in 2013–2017.

"Changes in survival rates over time varied by cancer type, with the largest survival improvements seen in prostate cancer, <u>kidney cancer</u>, multiple myeloma, non-Hodgkin lymphoma and tongue cancer," Mr Harvey said.

"While many cancers have high rates of survival, people diagnosed with cancers such as <u>pancreatic cancer</u>, <u>lung cancer</u> and mesothelioma have a less than 1 in 5 chance on average, of surviving at least five years after diagnosis."

Cancer <u>death</u> rates have also continued to drop since the 1980s.



"By the end of 2021, the cancer mortality rate is expected to reach a new low of 149 deaths per 100,000 people. It is expected that just under 50,000 people will die from cancer in 2021, an average of 135 deaths each day," Mr Harvey said.

While more males than females die from cancer, there has been a sharper decline in the cancer death rate for males than for females.

"The mortality rate for males reached a peak of 287 deaths per 100,000 in 1989 and the rate is estimated to have decreased by 37% to 182 deaths per 100,000 in 2021," Mr Harvey said.

"The overall decreasing trend for persons is largely due to the decline in the rate of death due to lung cancer, colorectal cancer, prostate cancer, cancer of unknown primary site and breast cancer."

These declines are likely due to a number of reasons. For example, smoking rates have declined dramatically since the 1960's; screening, early detection and monitoring programs (such as national bowel, breast and cervical cancer screening programs and PSA testing) have been introduced over the years; and therapeutic interventions have continued to improve.

The report also details information about rare and less common cancers, which generally have lower survival rates and are responsible for 2 in 5 cancer deaths.

While accounting for 30% of cancers diagnosed in 2017, rare and less common cancers accounted for close to 42% of deaths from cancer in that same year.

Rare cancers include bone cancer, mesothelioma, eye cancer and cancer of the nose and sinuses.



Stomach cancer, liver cancer, bladder cancer and brain cancer are among the group of less common cancers.

While overall progress has been made, the report shows that cancer remains a major cause of death in Australia.

"Cancers are responsible for more deaths than any other group of diseases, accounting for three in every 10 deaths in 2020," Mr. Harvey said.

"Lung cancer is expected to be the leading cause of death from cancer in 2021, followed by colorectal cancer, pancreatic cancer, <u>prostate cancer</u> and breast cancer."

The COVID-19 pandemic appears to have had at least some effect on the uptake of cancer-related services.

"After increasing by an average of 1% per year over the previous 20 years, the rate of cancer-related hospitalisations decreased by 1% between 2018-19 and 2019-20—noting that COVID-19 restrictions were in place only during the last quarter of 2019-20.

"COVID-19 restrictions also appear to have affected uptake of breast ultrasound, mammography, breast MRI and colonoscopy. For example, the number of people having Medicare Benefits Schedule-subsidized colonoscopies was 11% lower in 2020 compared with 2019, following average growth of around 3% per annum since 2011," Mr. Harvey said.

"The full impact of the COVID-19 pandemic on <u>cancer</u> diagnosis and treatment will not be known for some time."

Provided by Australian Institute of Health and Welfare (AIHW)



Citation: Cancer survival rates are improving in Australia (2021, November 30) retrieved 18 December 2023 from <u>https://medicalxpress.com/news/2021-11-cancer-survival-australia.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.