


# Lithium imaging method could shine new light on bipolar disorder, treatment

November 3 2021

---



Credit: CC0 Public Domain

Since 1949, lithium has been a mainstay for treating bipolar disorder (BD), a mental health condition marked by extreme mood swings. But scientists still don't have a clear understanding of how the drug works, or

why some patients respond better than others. Now, researchers reporting in *ACS Central Science* developed a method for imaging lithium in living cells, allowing them to discover that neurons from BD patients accumulate higher levels of lithium than healthy controls.

According to the National Institutes of Health, 4.4% of U.S. adults experience BD at some time in their lives. Studies have shown that lithium-based drugs can help stabilize mood and reduce suicide risk in people with BD. However, only about one-third of BD patients respond completely to lithium treatment, and the rest respond only partially or not at all. One reason could be that the drug has an extremely narrow therapeutic range: Below a certain blood serum level of lithium, most patients do not respond, but at a slightly higher level, they can experience [severe side effects](#). Being able to measure lithium concentrations directly in a patient's neurons could help scientists understand how lithium works as a drug, and then they could use this knowledge to optimize the dosage. So Yi Lu and colleagues wanted to develop a method to detect and measure lithium in living cells at therapeutically relevant concentrations.

The researchers used in vitro selection to identify a DNA enzyme (DNAzyme) that catalyzes the release of a fluorescent molecule from an RNA probe, thus producing a signal, only when lithium is present. The DNAzyme was 100 times more selective for lithium over other [metal ions](#), such as sodium and potassium, that are present at much higher concentrations in [human cells](#), and it was sensitive enough to detect lithium at concentrations within the therapeutic range. As a proof of concept, the researchers collected skin cells from BD patients and healthy donors, reprogrammed them to stem cells and then differentiated them into neurons. The team treated the neurons with the DNAzyme-based sensor and a therapeutically relevant dosage of lithium. Using [fluorescence microscopy](#), the researchers found that immature neurons from BD patients and healthy controls accumulated similar levels of

lithium, but mature neurons from BD patients accumulated higher levels of lithium than mature control neurons. The new lithium sensor is a powerful tool to better understand the effects of [lithium](#) in treating BD, the researchers say.

**More information:** DNAzyme-Based Lithium-Selective Imaging Reveals Higher Lithium Accumulation in Bipolar Disorder Patient-Derived Neurons, *ACS Central Science* (2021). [DOI: 10.1021/acscentsci.1c00843](#)

Provided by American Chemical Society

Citation: Lithium imaging method could shine new light on bipolar disorder, treatment (2021, November 3) retrieved 22 March 2023 from <https://medicalxpress.com/news/2021-11-lithium-imaging-method-bipolar-disorder.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--