


Researchers develop new imaging method to map alterations in the brain

February 23 2022



3D polarized light imaging scan of a mouse brain. The structural information about the orientation of the nerve fibres from this image served as a comparison for the new neutron measurements. Credit: Axer, Amunts et al. / INM-1, Forschungszentrum Jülich

The brain can be divided into so-called gray matter and white matter.


The white matter contains the axons that transmit stimuli. For faster stimulus transmission, the axons are wrapped in an insulating layer of myelin, in the same way as a cable is wrapped in rubber insulation to ensure that no electricity is lost along the way.

If the [myelin sheath](#) is damaged or degraded, this leads to impaired [brain](#) and body functions. In multiple sclerosis, for example, the insulating myelin layer is severely impaired. However, the exact causes of the disease are still unclear.

Researchers at the Jülich Institute of Neuroscience and Medicine and the Jülich Centre for Neutron Science have developed a new imaging method at the small-angle scattering facility KWS-1 at the MLZ to map the density, structure, and spatial orientation of nerve fibers and myelin.

New method complements previous procedures

"For a long time now, it has been possible to determine the spatial orientation of nerve fibers in brain slices using light microscopy and polarization," explains Dr. Heinrich Frielinghaus from the JCNS, head of the KWS-1 group, "but the exact structure and orientation of the myelin on the molecular scale cannot be detected using light." With the help of neutrons, the researchers can now for the first time determine both structure and orientation, and correlate these with polarization measurements.



Schematic diagram of the experimental setup at the small-angle scattering instrument. The neutron beam hits a brain section ("sample"). A detector records the scattering. Credit: Forschungszentrum Jülich

The special feature of this new method is that it allows scientists to image the [axons](#) bundled into [nerve fibers](#) at the same time as the insulating myelin sheath.

"With our neutron-based method, we can confirm the results of previous methods and provide additional information," says Frielinghaus.

Applications in medicine

"Since we work with brain slices, it is of course not a diagnostic procedure," explains Frielinghaus. With their new method, the researchers hope to better understand the causes of neurological diseases

in the future by being able to visualize structural changes in the brain more completely.

The resolution of the new method is still in the millimeter range. However, Frielinghaus and his colleagues expect that measurements precise to the micrometer will be possible in the future. This will enable them to study even the smallest structures in biological tissues.

More information: Santanu Maiti et al, Distribution and orientation of nerve fibers and myelin assembly in a brain section retrieved by small-angle neutron scattering, *Scientific Reports* (2021). [DOI: 10.1038/s41598-021-92995-2](https://doi.org/10.1038/s41598-021-92995-2)

Provided by Forschungszentrum Juelich

Citation: Researchers develop new imaging method to map alterations in the brain (2022, February 23) retrieved 30 March 2023 from <https://medicalxpress.com/news/2022-02-imaging-method-brain.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--