

SARS-CoV-2-linked hospitalization, ED encounters rare after Paxlovid

June 29 2022



Hospitalization and emergency department encounters related to severe



acute respiratory syndrome coronavirus 2 (SARS-CoV-2) occur rarely during the five to 15 days after Paxlovid treatment is dispensed for mild-to-moderate COVID-19, according to research published in the June 24 issue of the U.S. Centers for Disease Control and Prevention *Morbidity and Mortality Weekly Report*.

Deborah E. Malden, D.Phil., from the CDC in Atlanta, and colleagues used electronic health record data from Kaiser Permanente Southern California to describe <u>hospital admissions</u> and emergency department encounters related to SARS-CoV-2 infections during the five to 15 days after a five-day course of Paxlovid. Of the 5,287 persons aged ≥12 years who received Paxlovid during Dec. 31, 2021, to May 26, 2022, 73 and 8 percent had received three or more doses of COVID-19 vaccine and were unvaccinated, respectively.

The researchers identified six hospitalizations and 39 emergency department encounters considered related to SARS-CoV-2 infection during the five to 15 days after Paxlovid treatment was dispensed, representing

Citation: SARS-CoV-2-linked hospitalization, ED encounters rare after Paxlovid (2022, June 29) retrieved 1 April 2023 from

https://medicalxpress.com/news/2022-06-sars-cov-linked-hospitalization-ed-encounters-rare.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.