

Researchers discover neural plastic changes that improve functional recovery after cervical spinal cord injury

July 19 2022



Credit: CC0 Public Domain

Spinal cord and brain injury researchers at Indiana University School of Medicine are studying new alternatives to promote functional recovery after a spinal cord injury.

In a paper recently published in the *Journal of Clinical Investigation*, the team of researchers used models in the lab to investigate a unilateral spinal cord injury similar to Brown-Sequard Syndrome, a rare neurological condition where damage to the spinal cord in a person results in weakness or paralysis on one side of the body and a loss of sensation on the opposite side.

Wei Wu, Ph.D., assistant research professor of neurological surgery at IU School of Medicine, said the spinal cord injury model damaged the connection between the [left hemisphere](#) of the [brain](#) and the right side of the body, leading to a significant functional deficit to the right forelimb.

"The skilled function of upper limbs is very important for the quality of life in the patients with cervical spinal cord injury, but such functional recovery is very difficult to achieve in the [severe injury](#)," said Wu, first author of the paper and a member of the Spinal Cord and Brain Injury Research Group at Stark Neurosciences Research Institute. "We found that the intact corticospinal system in the opposite side of the brain and spinal cord can be modulated to at least partially take over the control of the forelimb that is damaged by the spinal cord injury, resulting in a forelimb functional improvement."

Each [hemisphere](#) in the brain controls the opposite side of the body, Wu added. Knowing this, the researchers discovered a spontaneous shift of the neural circuits after injury from the left hemisphere to the right. Although there are connections between the right hemisphere of the brain and the right side of the body through some relayed pathways after injury, Wu said that's not sufficient to support the motor recovery.

The team used an optogenetic neuromodulation strategy—stimulation using light—on the right hemisphere of the brain to modulate the motor cortex. This shifted additional neural circuits from the left side to the right side of the brain to dramatically increase and improve forelimb

function.

"New circuits in the whisker, jaw forelimb and neck areas in the right hemisphere of the brain are recruited to control the right forelimb," Wu said. "Interestingly, the beneficial neural plastic changes emerge both in the brain and the distal spinal cord after the optogenetic neuromodulation was applied on the motor cortex."

Wu said results of the study showed significant improvement to the forelimb; however, there are still many challenges ahead, since the complete digital recovery is not succeeded.

The research team will continue to understand and mediate this transhemispheric neural reorganization to further improve the [functional recovery](#) after the spinal cord injury, Wu said. He hopes the discoveries from this study will be translated to a clinical therapeutic strategy for patients with [spinal cord](#) injuries.

More information: Wei Wu et al, Transhemispheric cortex remodeling promotes forelimb recovery after spinal cord injury, *Journal of Clinical Investigation* (2022). [DOI: 10.1172/jci.insight.158150](https://doi.org/10.1172/jci.insight.158150)

Provided by Indiana University School of Medicine

Citation: Researchers discover neural plastic changes that improve functional recovery after cervical spinal cord injury (2022, July 19) retrieved 30 January 2023 from <https://medicalxpress.com/news/2022-07-neural-plastic-functional-recovery-cervical.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.