


# Particle radioactivity linked to pollution-associated heart attack and stroke death

October 5 2022

---



Credit: CC0 Public Domain

Particle radioactivity, a characteristic of air pollution that reflects the colorless, odorless gas radon found in fine particulate matter (PM<sub>2.5</sub>) air

pollution, enhances PM<sub>2.5</sub> toxicity and increases risk of death from cardiovascular disease, especially from heart attack or stroke, according to new research published today in the *Journal of the American Heart Association*.

Previous scientific research has confirmed that PM<sub>2.5</sub>, a component of air pollution, causes cardiovascular [disease](#) and [death](#), and that PM<sub>2.5</sub> exposure is a modifiable cardiovascular disease risk factor. In 2021, the [American Heart Association](#) joined with three other leading cardiovascular organizations urging the [medical community](#) and [health authorities](#) to mitigate the impact of air pollution on people's health. According to the statement, an estimated 6.7 million deaths in 2019, or 12 percent of all deaths worldwide, were attributable to outdoor or household air pollution. As many as half of these were due to cardiovascular disease. Air pollution also increases the risk of [heart attack](#), stroke, diabetes and [respiratory diseases](#).

Particle radioactivity is a characteristic of particulate matter that reflects radon, which primarily comes from radon gas, a radioactive, colorless and odorless gas. The particle radioactivity occurs naturally as a product of radioactive decay of uranium found in soil and rocks. Radon migrates into the atmosphere, decaying to alpha-, beta- and gamma-radiation-emitting isotopes.

"We know that PM<sub>2.5</sub> are very small particles in the air that can be inhaled and cause many health problems. However, little is known about which physical, chemical or biological properties of PM<sub>2.5</sub> fuel its toxicity," said study author Shuxin Dong, S.M., a Ph.D. student in population health sciences at Harvard T.H. Chan School of Public Health in Boston. "We studied gross beta-activity, a property of [fine particulate matter](#) that is a result of radon that attaches to particles and makes them radioactive, resulting in particle radioactivity. When inhaled, these very [small particles](#) penetrate deeply into the lungs and enter the bloodstream

and circulate throughout the body."

The researchers used spatiotemporal predictions of gross beta-activity, a way to use different variables across space and time, to provide refined predictions of exposure. Examining health records from more than 700,000 non-accidental deaths in Massachusetts between 2001 and 2015, they estimated how long-term (months/year) gross beta-activity exposure impacts death from cardiovascular disease, heart attack or stroke and death from all non-accidental causes. They also predicted  $PM_{2.5}$  on cardiovascular disease-related death and examined the interaction between  $PM_{2.5}$  and particle radioactivity.

The study found:

- Chronic particle radioactivity and  $PM_{2.5}$  exposure were similarly associated with increased risks of death from total cardiovascular disease, heart attack or stroke and all causes of non-accidental death.
- Based on the middle 50% of the data spread, particle radioactivity exposure alone was associated with a 16% increased risk of death from heart attack; an 11% increased risk of death from stroke; a 7% increased risk of death from all types of cardiovascular disease; and a 4% increased risk of death from all non-accidental causes.
- Based on the middle 50% of the data spread,  $PM_{2.5}$  exposure alone increased the risk of death from heart attack by 6%; death from stroke by 11%; death from all cardiovascular disease by 12%; and death from all non-accidental causes by 10%.

"The risk of death from cardiovascular disease, [heart attack](#) or stroke and all causes due to  $PM_{2.5}$  was higher and, therefore, more toxic when gross beta-activity levels were higher," Dong said. "These findings suggest that particle radioactivity increases the risk of death from

[cardiovascular disease](#) and enhances the damage from particulate matter. This must be further investigated and may lead to targeted, cost-effective air quality regulations."

A study limitation is that the research was based on information from one state, Massachusetts, and therefore, the results may not be generalizable to the rest of the U.S.

Co-authors are Petros Koutrakis, Ph.D.; Longxiang Li, Ph.D.; Brent A. Coull, Ph.D.; Joel Schwartz, Ph.D.; Anna Kosheleva, M.S.; and Antonella Zanobetti, Ph.D. Authors' disclosures are listed in the manuscript.

**More information:** Synergistic Effects of Particle Radioactivity (Gross  $\beta$  Activity) and Particulate Matter  $\leq 2.5\mu\text{m}$  Aerodynamic Diameter on Cardiovascular Disease Mortality, *Journal of the American Heart Association* (2022). [DOI: 10.1161/JAHA.121.025470](https://doi.org/10.1161/JAHA.121.025470)

Provided by American Heart Association

Citation: Particle radioactivity linked to pollution-associated heart attack and stroke death (2022, October 5) retrieved 4 February 2024 from <https://medicalxpress.com/news/2022-10-particle-radioactivity-linked-pollution-associated-heart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.