


# Is metabolic syndrome a modifiable risk factor for gout?

November 23 2022

---



Credit: Pixabay/CC0 Public Domain

In a study published in *Arthritis & Rheumatology* that included nearly 1.3 million men aged 20–39 years who participated in three serial health check-ups at two-year intervals, men with metabolic syndrome (MetS) and those who developed MetS—especially those with the MetS components of elevated triglycerides and abdominal obesity—had higher

risks of developing gout.

Among participants, 18,473 developed [gout](#), and those with MetS at all checkups had a nearly four-fold higher risk than participants who were MetS-free. Development of MetS more than doubled the risk of incident gout, whereas recovery from MetS reduced incident gout risk by nearly half.

"This is the first large-scale study to explore the association between dynamic changes in MetS and risk of gout," said co-corresponding author Jaejoon Lee, MD, Ph.D. of the Sungkyunkwan University School of Medicine, in South Korea. "Prevention and recovery from MetS can significantly lower the risk of gout in young adults."

**More information:** Yeonghee Eun et al, Altered Risk of Incident Gout According to Changes in Metabolic Syndrome Status: A Nationwide Population-Based Cohort Study of 1.29 Million Young Men, *Arthritis & Rheumatology* (2022). [DOI: 10.1002/art.42381](https://doi.org/10.1002/art.42381)

Provided by Wiley

Citation: Is metabolic syndrome a modifiable risk factor for gout? (2022, November 23) retrieved 29 April 2023 from <https://medicalxpress.com/news/2022-11-metabolic-syndrome-factor-gout.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--